

RNF126 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21884c**Specification**

RNF126 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9BV68
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	33861

RNF126 Antibody (Center) - Additional Information**Gene ID** 55658**Other Names**

E3 ubiquitin-protein ligase RNF126, 632-, RING finger protein 126
{ECO:0000312|HGNC:HGNC:21151}, RNF126 (<a
href="http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=21151"
target="_blank">HGNC:21151)

Target/Specificity

This RNF126 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 84-115 amino acids from the Central region of human RNF126.

Dilution

WB~~1:2000
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RNF126 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

RNF126 Antibody (Center) - Protein Information**Name** RNF126 {ECO:0000303|PubMed:23026136, ECO:0000312|HGNC:HGNC:21151}

Function E3 ubiquitin-protein ligase that mediates ubiquitination of target proteins (PubMed:[23277564](#), PubMed:[24275455](#), PubMed:[24981174](#), PubMed:[36563124](#)). Depending on the associated E2 ligase, mediates 'Lys-27'-, 'Lys-29'-, 'Lys-48'- and/or 'Lys-63'-linked polyubiquitination of substrates (PubMed:[36563124](#)). Part of a BAG6-dependent quality control process ensuring that proteins of the secretory pathway that are mislocalized to the cytosol are degraded by the proteasome. Probably acts by providing the ubiquitin ligase activity associated with the BAG6 complex and be responsible for ubiquitination of the hydrophobic mislocalized proteins and their targeting to the proteasome (PubMed:[24981174](#), PubMed:[29042515](#)). May also play a role in the endosomal recycling of IGF2R, the cation-independent mannose-6-phosphate receptor (PubMed:[24275455](#)). May play a role in the endosomal sorting and degradation of several membrane receptors including EGFR, FLT3, MET and CXCR4, by mediating their ubiquitination (PubMed:[23418353](#)). By ubiquitinating CDKN1A/p21 and targeting it for degradation, may also promote cell proliferation (PubMed:[23026136](#)). May monoubiquitinate AICDA (PubMed:[23277564](#)). Acts as a regulator of DNA repair by mediating 'Lys-27'- and 'Lys-29'-linked polyubiquitination of MRE11, thereby promoting the exonuclease activity of MRE11 (PubMed:[36563124](#)).

Cellular Location

Cytoplasm. Nucleus


Tissue Location

Highly expressed in liver and testis.

RNF126 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RNF126 Antibody (Center) - Images

Anti-RNF126 Antibody (Center) at 1:2000 dilution + CEM whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

RNF126 Antibody (Center) - Background

E3 ubiquitin-protein ligase that regulates several biological processes through ubiquitination of various target proteins. Depending on the associated E2 ligase, mediates 'Lys- 48'- and 'Lys-63'-linked polyubiquitination of substrates. Through their polyubiquitination, may play a role in the endosomal sorting and degradation of several membrane receptors including EGFR, FLT3, MET and CXCR4. May also be part of a BAG6-dependent quality control process ensuring that proteins of the secretory pathway that are mislocalized to the cytosol are degraded by the proteasome. May provide the ubiquitin ligase activity associated with the BAG6 complex and be responsible for ubiquitination of the mislocalized proteins and their targeting to the proteasome (PubMed:24981174). May also play a role in the endosomal recycling of IGF2R, the cation-independent mannose-6-phosphate receptor (PubMed:24275455). By ubiquitinating CDKN1A/p21 and targeting it for degradation, may also promote cell proliferation (PubMed:23026136). May monoubiquitinate AICDA (PubMed:23277564).

RNF126 Antibody (Center) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Daub H.,et al.Mol. Cell 31:438-448(2008).
Olsen J.V.,et al.Sci. Signal. 3:RA3-RA3(2010).
Zhi X.,et al.Cancer Res. 73:385-394(2013).
Smith C.J.,et al.J. Cell Sci. 126:1366-1380(2013).