

TFDP1 Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21890a**Specification**

TFDP1 Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	Q14186
Other Accession	Q17QZ4 , Q08639
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	45070

TFDP1 Antibody (N-Term) - Additional Information**Gene ID** 7027**Other Names**

Transcription factor Dp-1, DRTF1-polypeptide 1, DRTF1, E2F dimerization partner 1, TFDP1, DP1

Target/Specificity

This TFDP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 83-114 amino acids from human TFDP1.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TFDP1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

TFDP1 Antibody (N-Term) - Protein Information**Name** TFDP1**Synonyms** DP1

Function Can stimulate E2F-dependent transcription. Binds DNA cooperatively with E2F family members through the E2 recognition site, 5'-TTTC[CG]CGC-3', found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication (PubMed:[7739537](#), PubMed:[8405995](#)). The E2F1:DP complex appears to mediate both cell proliferation and apoptosis. Blocks adipocyte differentiation by repressing CEBPA binding to its target gene promoters (PubMed:[20176812](#)).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q08639}. Cytoplasm {ECO:0000250|UniProtKB:Q08639}. Note=Shuttles between the cytoplasm and nucleus and translocates into the nuclear compartment upon heterodimerization with E2F1. {ECO:0000250|UniProtKB:Q08639}

Tissue Location


Highest levels in muscle. Also expressed in brain, placenta, liver and kidney. Lower levels in lung and pancreas. Not detected in heart

TFDP1 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TFDP1 Antibody (N-Term) - Images



All lanes : Anti-TFDP1 Antibody (N-Term) at 1:2000 dilution Lane 1: C2C12 whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

TFDP1 Antibody (N-Term) - Background

Can stimulate E2F-dependent transcription. Binds DNA cooperatively with E2F family members through the E2 recognition site, 5'-TTTC[CG]CGC-3', found in the promoter region of a number of genes whose products are involved in cell cycle regulation or in DNA replication. The DP2/E2F complex functions in the control of cell-cycle progression from G1 to S phase. The E2F1/DP complex appears to mediate both cell proliferation and apoptosis.

TFDP1 Antibody (N-Term) - References

Helin K.,et al.Genes Dev. 7:1850-1861(1993).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham A.,et al.Nature 428:522-528(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Bandara L.R.,et al.EMBO J. 13:3104-3114(1994).