

**GPR124 Antibody (C-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21894b****Specification**

---

**GPR124 Antibody (C-Term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q96PE1</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	142647

**GPR124 Antibody (C-Term) - Additional Information****Gene ID** 25960**Other Names**

G-protein coupled receptor 124, Tumor endothelial marker 5, GPR124, KIAA1531, TEM5

**Target/Specificity**

This GPR124 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1259-1293 amino acids from human GPR124.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GPR124 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**GPR124 Antibody (C-Term) - Protein Information****Name** ADGRA2 ([HGNC:17849](#))**Function** Endothelial receptor which functions together with RECK to enable brain endothelial cells to selectively respond to Wnt7 signals (WNT7A or WNT7B) (PubMed:[28289266](#), PubMed:[30026314](#)). Plays a key role in Wnt7-specific responses, such as endothelial cell sprouting

and migration in the forebrain and neural tube, and establishment of the blood-brain barrier (By similarity). Acts as a Wnt7-specific coactivator of canonical Wnt signaling: required to deliver RECK-bound Wnt7 to frizzled by assembling a higher-order RECK-ADGRA2-Fzd-LRP5-LRP6 complex (PubMed:[30026314](#)). ADGRA2-tethering function does not rely on its G-protein coupled receptor (GPCR) structure but instead on its combined capacity to interact with RECK extracellularly and recruit the Dishevelled scaffolding protein intracellularly (PubMed:[30026314](#)). Binds to the glycosaminoglycans heparin, heparin sulfate, chondroitin sulfate and dermatan sulfate (PubMed:[16982628](#)).

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, filopodium. Note=Enriched at lateral cell borders and also at sites of cell-ECM (extracellular matrix) contact

#### Tissue Location


Expressed in endothelial cells (at protein level) (PubMed:15021905, PubMed:16982628). Abundantly expressed in heart, placenta, ovary, small intestine, and colon (PubMed:15021905)

### GPR124 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GPR124 Antibody (C-Term) - Images



Anti-GPR124 Antibody (C-Term) at 1:2000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 143 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### GPR124 Antibody (C-Term) - Background

Endothelial receptor that acts as an essential regulator of CNS angiogenesis. Required for normal endothelial cell sprouting and migration in the forebrain and neural tube (By similarity).

#### **GPR124 Antibody (C-Term) - References**

Carson-Walter E.B.,et al.Cancer Res. 61:6649-6655(2001).  
Nagase T.,et al.DNA Res. 7:143-150(2000).  
Nakajima D.,et al.DNA Res. 9:99-106(2002).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Nusbaum C.,et al.Nature 439:331-335(2006).