

C18orf32 Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21952b**Specification**

C18orf32 Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	Q8TCD1
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	8669

C18orf32 Antibody (C-Term) - Additional Information**Gene ID** 497661**Other Names**

UPF0729 protein C18orf32, Putative NF-kappa-B-activating protein 200, C18orf32

Target/Specificity

This C18orf32 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 26-58 amino acids from human C18orf32.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C18orf32 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

C18orf32 Antibody (C-Term) - Protein Information**Name** C18orf32**Function** May activate the NF-kappa-B signaling pathway.**Cellular Location**


Endoplasmic reticulum. Lipid droplet

C18orf32 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C18orf32 Antibody (C-Term) - Images



All lanes : Anti-C18orf32 Antibody (C-Term) at 1:2000 dilution Lane 1: human brain lysate Lane 2: LNCaP whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 9 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

C18orf32 Antibody (C-Term) - Background

May activate the NF-kappa-B signaling pathway.

C18orf32 Antibody (C-Term) - References

Matsuda A.,et al.Oncogene 22:3307-3318(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).