

**TACC3 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21957c****Specification**

---

**TACC3 Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q9Y6A5</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	90360

**TACC3 Antibody (Center) - Additional Information****Gene ID** 10460**Other Names**

Transforming acidic coiled-coil-containing protein 3, ERIC-1, TACC3, ERIC1

**Target/Specificity**

This TACC3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 317-350 amino acids from the Central region of human TACC3.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TACC3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**TACC3 Antibody (Center) - Protein Information****Name** TACC3**Synonyms** ERIC1**Function** Plays a role in the microtubule-dependent coupling of the nucleus and the centrosome.

Involved in the processes that regulate centrosome-mediated interkinetic nuclear migration (INM) of neural progenitors (By similarity). Acts as a component of the TACC3/ch-TOG/clathrin complex proposed to contribute to stabilization of kinetochore fibers of the mitotic spindle by acting as inter-microtubule bridge. The TACC3/ch-TOG/clathrin complex is required for the maintenance of kinetochore fiber tension (PubMed:[21297582](#), PubMed:[23532825](#)). May be involved in the control of cell growth and differentiation. May contribute to cancer (PubMed:[14767476](#)).

#### Cellular Location


Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole {ECO:0000250|UniProtKB:Q9PTG8}. Note=In complex with CKAP5 localized to microtubule plus-ends in mitosis and interphase. In complex with CKAP5 and clathrin localized to inter-microtubule bridges in mitotic spindles.

#### TACC3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TACC3 Antibody (Center) - Images



All lanes : Anti-TACC3 Antibody (Center) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: HCT116 whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: K562 whole cell lysate Lane 5: U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 90 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

#### TACC3 Antibody (Center) - Background

Plays a role in the microtubule-dependent coupling of the nucleus and the centrosome. Involved in

the processes that regulate centrosome-mediated interkinetic nuclear migration (INM) of neural progenitors (By similarity). May be involved in the control of cell growth and differentiation. May contribute to cancer.

#### **TACC3 Antibody (Center) - References**

Still I.H.,et al.Genomics 58:165-170(1999).  
McKeveney P.J.,et al.Br. J. Haematol. 112:1016-1024(2001).  
Gangisetty O.,et al.Oncogene 23:2559-2563(2004).  
Beausoleil S.A.,et al.Nat. Biotechnol. 24:1285-1292(2006).  
Cantin G.T.,et al.J. Proteome Res. 7:1346-1351(2008).