

**PRAM1 Antibody (C-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22018b**

**Specification**

---

**PRAM1 Antibody (C-Term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q96QH2</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	73969

**PRAM1 Antibody (C-Term) - Additional Information**

**Gene ID** 84106

**Other Names**

PML-RARA-regulated adapter molecule 1, PRAM, PRAM-1, PRAM1

**Target/Specificity**

This PRAM1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 587-618 amino acids from human PRAM1.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PRAM1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PRAM1 Antibody (C-Term) - Protein Information**

**Name** PRAM1

**Function** May be involved in myeloid differentiation. May be involved in integrin signaling in neutrophils. Binds to PtdIns(4)P.

**Tissue Location**


Expressed in peripheral blood leukocytes and bone marrow. Expressed in monocytes, and to a lesser extent in granulocytes and lymphocytes. Not expressed in non hematopoietic tissues except in lung.

### **PRAM1 Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **PRAM1 Antibody (C-Term) - Images**



Anti-PRAM1 Antibody (C-Term) at 1:2000 dilution + THP-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 79 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### **PRAM1 Antibody (C-Term) - Background**

May be involved in myeloid differentiation. May be involved in integrin signaling in neutrophils. Binds to PtdIns(4)P.

### **PRAM1 Antibody (C-Term) - References**

Moog-Lutz C., et al. J. Biol. Chem. 276:22375-22381(2001).  
Grimwood J., et al. Nature 428:529-535(2004).  
Denis F.M., et al. J. Biol. Chem. 280:9043-9048(2005).  
Heuer K., et al. J. Mol. Biol. 361:94-104(2006).  
Choudhary C., et al. Science 325:834-840(2009).