

JMJD1C Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22062b**Specification**

JMJD1C Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	Q15652
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	284525

JMJD1C Antibody (C-Term) - Additional Information**Gene ID** 221037**Other Names**

Probable JmjC domain-containing histone demethylation protein 2C, 1.14.11.-, Jumonji domain-containing protein 1C, Thyroid receptor-interacting protein 8, TR-interacting protein 8, TRIP-8, JMJD1C, JHDM2C, KIAA1380, TRIP8

Target/Specificity

This JMJD1C antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 2019-2051 amino acids from human JMJD1C.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

JMJD1C Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

JMJD1C Antibody (C-Term) - Protein Information**Name** JMJD1C**Synonyms** JHDM2C, KIAA1380, TRIP8

Function Probable histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Demethylation of Lys residue generates formaldehyde and succinate. May be involved in hormone-dependent transcriptional activation, by participating in recruitment to androgen-receptor target genes (By similarity).

Cellular Location


Nucleus.

JMJD1C Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

JMJD1C Antibody (C-Term) - Images



Anti-JMJD1C Antibody (C-Term) at 1:1000 dilution + human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 285 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

JMJD1C Antibody (C-Term) - Background

Probable histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Demethylation of Lys residue generates formaldehyde and succinate. May be involved in hormone-dependent transcriptional activation, by participating in recruitment to androgen-receptor target genes (By similarity).

JMJD1C Antibody (C-Term) - References

Wolf S.S., et al. Arch. Biochem. Biophys. 460:56-66(2007).

Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Deloukas P.,et al.Nature 429:375-381(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Nagase T.,et al.DNA Res. 7:65-73(2000).