

LSR Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22086a

Specification

LSR Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	Q86X29
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	71439

LSR Antibody (N-Term) - Additional Information

Gene ID 51599

Other Names

Lipolysis-stimulated lipoprotein receptor, LSR, LISCH

Target/Specificity

This LSR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 10-43 amino acids from human LSR.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LSR Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

LSR Antibody (N-Term) - Protein Information

Name LSR ([HGNC:29572](#))

Synonyms ILDR3, LISCH

Function Probable role in the clearance of triglyceride-rich lipoprotein from blood. Binds

chylomicrons, LDL and VLDL in presence of free fatty acids and allows their subsequent uptake in the cells (By similarity). Maintains epithelial barrier function by recruiting MARVELD2/tricellulin to tricellular tight junctions (By similarity).

Cellular Location


Cell membrane; Single-pass type I membrane protein. Cell junction, tight junction Note=Located at tricellular contacts.

LSR Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LSR Antibody (N-Term) - Images



Anti-LSR Antibody (N-Term) at 1:2000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 71 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

LSR Antibody (N-Term) - Background

Probable role in the clearance of triglyceride-rich lipoprotein from blood. Binds chylomicrons, LDL and VLDL in presence of free fatty acids and allows their subsequent uptake in the cells (By similarity).

LSR Antibody (N-Term) - References

Teramoto T., et al. Submitted (FEB-1999) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).

Otsuki T., et al. DNA Res. 12:117-126(2005).
Grimwood J., et al. Nature 428:529-535(2004).
Tang L.-Y., et al. Mol. Cell. Proteomics 6:1952-1967(2007).