

ALG6 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22095c**Specification**

ALG6 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O9Y672
Other Accession	O3TAE8 , O5NVS8 , O3T1L5
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	58121

ALG6 Antibody (Center) - Additional Information**Gene ID** 29929**Other Names**

Dolichyl pyrophosphate Man9GlcNAc2 alpha-1, 3-glucosyltransferase, 2.4.1.267, Asparagine-linked glycosylation protein 6 homolog, Dol-P-Glc:Man(9)GlcNAc(2)-PP-Dol alpha-1, 3-glucosyltransferase, Dolichyl-P-Glc:Man9GlcNAc2-PP-dolichyl glucosyltransferase, ALG6

Target/Specificity

This ALG6 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 241-274 amino acids from the Central region of human ALG6.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ALG6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ALG6 Antibody (Center) - Protein Information**Name** ALG6 ([HGNC:23157](#))

Function Dolichyl pyrophosphate Man9GlcNAc2 alpha-1,3- glucosyltransferase that operates in the biosynthetic pathway of dolichol-linked oligosaccharides, the glycan precursors employed in protein asparagine (N)-glycosylation. The assembly of dolichol-linked oligosaccharides begins on the cytosolic side of the endoplasmic reticulum membrane and finishes in its lumen. The sequential addition of sugars to dolichol pyrophosphate produces dolichol-linked oligosaccharides containing fourteen sugars, including two GlcNAcs, nine mannoses and three glucoses. Once assembled, the oligosaccharide is transferred from the lipid to nascent proteins by oligosaccharyltransferases. In the lumen of the endoplasmic reticulum, adds the first glucose residue from dolichyl phosphate glucose (Dol-P- Glc) onto the lipid-linked oligosaccharide intermediate Man(9)GlcNAc(2)-PP-Dol to produce Glc(1)Man(9)GlcNAc(2)-PP-Dol. Glc(1)Man(9)GlcNAc(2)-PP-Dol is a substrate for ALG8, the following enzyme in the biosynthetic pathway.

Cellular Location


Endoplasmic reticulum membrane; Multi-pass membrane protein

ALG6 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ALG6 Antibody (Center) - Images



All lanes : Anti-ALG6 Antibody (Center) at 1:2000 dilution Lane 1: human fetal lung lysate Lane 2: human fetal heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

ALG6 Antibody (Center) - Background

Adds the first glucose residue to the lipid-linked oligosaccharide precursor for N-linked glycosylation. Transfers glucose from dolichyl phosphate glucose (Dol-P-Glc) onto the lipid-linked oligosaccharide Man(9)GlcNAc(2)-PP-Dol.

ALG6 Antibody (Center) - References

Imbach T.,et al.Proc. Natl. Acad. Sci. U.S.A. 96:6982-6987(1999).
Mao Y.M.,et al.Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.