

Profilin-1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22110a**Specification**

Profilin-1 Antibody - Product Information

Application	WB, FC,E
Primary Accession	P07737
Other Accession	P02584 , P62962 , P62963
Reactivity	Human, Mouse, Rat
Predicted	Bovine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	15054

Profilin-1 Antibody - Additional Information**Gene ID** 5216**Other Names**

Profilin-1, Epididymis tissue protein Li 184a, Profilin I, PFN1

Target/Specificity

This Profilin-1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 108-140 amino acids from the human region of human Profilin-1.

Dilution

WB~~1:2000

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Profilin-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Profilin-1 Antibody - Protein Information**Name** PFN1**Function** Binds to actin and affects the structure of the cytoskeleton. At high concentrations,

profilin prevents the polymerization of actin, whereas it enhances it at low concentrations. By binding to PIP2, it inhibits the formation of IP3 and DG. Inhibits androgen receptor (AR) and HTT aggregation and binding of G-actin is essential for its inhibition of AR.

Cellular Location

Cytoplasm, cytoskeleton.

Tissue Location


Expressed in epididymis (at protein level).

Profilin-1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Profilin-1 Antibody - Images



All lanes : Anti-Profilin-1 Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: HUVEC whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: 293 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lane 6: C6 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 15 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing NIH/3T3 cells stained with AP22110a (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22110a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.



Overlay histogram showing HeLa cells stained with AP22110a (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22110a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

Profilin-1 Antibody - Background

Binds to actin and affects the structure of the cytoskeleton. At high concentrations, profilin prevents the polymerization of actin, whereas it enhances it at low concentrations. By binding to PIP2, it inhibits the formation of IP3 and DG. Inhibits androgen receptor (AR) and HTT aggregation and binding of G-actin is essential for its inhibition of AR.

Profilin-1 Antibody - References

Kwiatkowski D.J., et al. J. Biol. Chem. 263:5910-5915(1988).

Li J.,et al.Mol. Cell. Proteomics 9:2517-2528(2010).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.