

UHRF1BP1 Antibody (C-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22184b**Specification**

UHRF1BP1 Antibody (C-Term) - Product Information

Application	WB,E
Primary Accession	Q6BDS2
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	159485

UHRF1BP1 Antibody (C-Term) - Additional Information**Gene ID** 54887**Other Names**

UHRF1-binding protein 1, ICBP90-binding protein 1, Ubiquitin-like containing PHD and RING finger domains 1-binding protein 1, UHRF1BP1, C6orf107

Target/Specificity

This UHRF1BP1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1096-1129 amino acids from the human UHRF1BP1.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UHRF1BP1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

UHRF1BP1 Antibody (C-Term) - Protein Information**Name** BLTP3A ([HGNC:21216](#))**Synonyms** C6orf107, UHRF1BP1**Function** Tube-forming lipid transport protein which probably mediates the transfer of lipids

between membranes at organelle contact sites (PubMed:[35499567](#)). May be involved in the retrograde traffic of vesicle clusters in the endocytic pathway to the Golgi complex (PubMed:[35499567](#)).


Cellular Location

Late endosome.

UHRF1BP1 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UHRF1BP1 Antibody (C-Term) - Images

Anti-UHRF1BP1 Antibody (C-Term) at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 159 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

UHRF1BP1 Antibody (C-Term) - Background

May act as a negative regulator of cell growth.

UHRF1BP1 Antibody (C-Term) - References

Unoki M.,et al.Oncogene 23:7601-7610(2004).
Mungall A.J.,et al.Nature 425:805-811(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).
Gauci S.,et al.Anal. Chem. 81:4493-4501(2009).

