

MUT Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22354a**Specification**

MUT Antibody (N-Term) - Product Information

Application	WB, IF, IHC-P-Leica,E
Primary Accession	P22033
Other Accession	Q9GK13 , Q8HXX1 , P16332 , Q5RFN2
Reactivity	Human, Mouse
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG

MUT Antibody (N-Term) - Additional Information**Gene ID** 4594**Other Names**

Methylmalonyl-CoA mutase, mitochondrial, MCM, 5.4.99.2, Methylmalonyl-CoA isomerase, MUT

Target/Specificity

This MUT antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-66 amino acids from the human region of human MUT.

Dilution

WB~~1:2000
IF~~1:50~200
IHC-P-Leica~~1:250
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MUT Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

MUT Antibody (N-Term) - Protein Information**Name** MMUT ([HGNC:7526](#))

Function Catalyzes the reversible isomerization of methylmalonyl-CoA (MMCoA) (generated from branched-chain amino acid metabolism and degradation of dietary odd chain fatty acids and cholesterol) to succinyl-CoA (3-carboxypropionyl-CoA), a key intermediate of the tricarboxylic acid cycle.

Cellular Location


Mitochondrion matrix. Mitochondrion. Cytoplasm

MUT Antibody (N-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MUT Antibody (N-Term) - Images



All lanes : Anti-MUT Antibody (N-Term) at 1:2000 dilution Lane 1: Human liver lysate Lane 2: Mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-MUT Antibody (N-Term) at 1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded human testis tissue using AP22354a performed on the Leica® BOND RXm. Samples were incubated with primary antibody(1/250) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

MUT Antibody (N-Term) - Background

Involved in the degradation of several amino acids, odd- chain fatty acids and cholesterol via propionyl-CoA to the tricarboxylic acid cycle. MCM has different functions in other species.

MUT Antibody (N-Term) - References

- Jansen R.,et al.Genomics 4:198-205(1989).
- Nham S.U.,et al.Genomics 8:710-716(1990).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
- Mungall A.J.,et al.Nature 425:805-811(2003).

