

IFT57 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22359c**Specification**

IFT57 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q9NWB7
Other Accession	Q5EA95
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	49108

IFT57 Antibody (Center) - Additional Information**Gene ID** 55081**Other Names**

Intraflagellar transport protein 57 homolog, Dermal papilla-derived protein 8, Estrogen-related receptor beta-like protein 1, HIP1-interacting protein, MHS4R2, IFT57, DERP8, ESRRBL1, HIPPI

Target/Specificity

This IFT57 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 159-193 amino acids from the Central region of human IFT57.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IFT57 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

IFT57 Antibody (Center) - Protein Information**Name** IFT57

Synonyms DERP8, ESRRBL1, HIPPI

Function Required for the formation of cilia. Plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway (By similarity). Has pro-apoptotic function via its interaction with HIP1, leading to recruit caspase-8 (CASP8) and trigger apoptosis. Has the ability to bind DNA sequence motif 5'- AAAGACATG-3' present in the promoter of caspase genes such as CASP1, CASP8 and CASP10, suggesting that it may act as a transcription regulator; however the relevance of such function remains unclear.

Cellular Location

Cell projection, cilium {ECO:0000250|UniProtKB:Q8BXG3}. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q5EA95}. Note=Concentrates within the inner segment of cilia.

Tissue Location


Present in many tissues such as brain, thymus, lymph node, lung, liver, skin and kidney (at protein level)

IFT57 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IFT57 Antibody (Center) - Images



All lanes : Anti-IFT57 Antibody (Center) at 1:2000 dilution Lane 1: Human brain lysate Lane 2: Human lung lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

IFT57 Antibody (Center) - Background

Required for the formation of cilia. Plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway (By similarity). Has pro- apoptotic function via its interaction with HIP1, leading to recruit caspase-8 (CASP8) and trigger apoptosis. Has the ability to bind DNA sequence motif 5'-AAAGACATG-3' present in the promoter of caspase genes such as CASP1, CASP8 and CASP10, suggesting that it may act as a transcription regulator; however the relevance of such function remains unclear.

IFT57 Antibody (Center) - References

Gervais F.G.,et al.Nat. Cell Biol. 4:95-105(2002).
Pasutto F.,et al.Submitted (MAR-1999) to the EMBL/GenBank/DDBJ databases.
Ikeda A.,et al.Submitted (MAY-1998) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.