

P21CIP1(T57) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22485a**Specification**

P21CIP1(T57) Antibody - Product Information

Application	WB,E
Primary Accession	P38936
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	18119

P21CIP1(T57) Antibody - Additional Information**Gene ID** 1026**Other Names**

Cyclin-dependent kinase inhibitor 1, CDK-interacting protein 1, Melanoma differentiation-associated protein 6, MDA-6, p21, CDKN1A ([HGNC:1784](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=1784))

Target/Specificity

This P21CIP1(T57) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from the human region of human P21CIP1(T57).

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

P21CIP1(T57) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

P21CIP1(T57) Antibody - Protein Information**Name** CDKN1A ([HGNC:1784](#))**Function** Plays an important role in controlling cell cycle progression and DNA damage-induced

G2 arrest (PubMed:[9106657](#)). Involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Also involved in p53-independent DNA damage-induced G2 arrest mediated by CREB3L1 in astrocytes and osteoblasts (By similarity). Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:[11595739](#)). Negatively regulates the CDK4- and CDK6-driven phosphorylation of RB1 in keratinocytes, thereby resulting in the release of E2F1 and subsequent transcription of E2F1-driven G1/S phase promoting genes (By similarity).

Cellular Location

Cytoplasm. Nucleus


Tissue Location

Expressed in all adult tissues, with 5-fold lower levels observed in the brain

P21CIP1(T57) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

P21CIP1(T57) Antibody - Images

All lanes: Anti-P21CIP1(T57) Antibody at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 22 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

P21CIP1(T57) Antibody - Background

Plays an important role in controlling cell cycle progression and DNA damage-induced G2 arrest (PubMed:9106657). Involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Also involved in p53-independent DNA damage-induced G2 arrest mediated by CREB3L1 in astrocytes and osteoblasts (By similarity). Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:11595739). Negatively regulates the CDK4- and CDK6-driven phosphorylation of RB1 in keratinocytes, thereby resulting in the release of E2F1 and subsequent transcription of E2F1-driven G1/S phase promoting genes (By similarity).

P21CIP1(T57) Antibody - References

Harper J.W.,et al.Cell 75:805-816(1993).
El-Deiry W.S.,et al.Cell 75:817-825(1993).
Xiong Y.,et al.Nature 366:701-704(1993).
Jiang H.,et al.Mol. Cell. Differ. 1:285-299(1993).
Jiang H.,et al.Oncogene 10:1855-1864(1995).