

ART1 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2311a**Specification**

ART1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P52961
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36335
Antigen Region	14-43

ART1 Antibody (N-term) - Additional Information**Gene ID** 417**Other Names**

GPI-linked NAD(P)(+)-arginine ADP-ribosyltransferase 1, ADP-ribosyltransferase C2 and C3 toxin-like 1, ARTC1, Mono(ADP-ribosyl)transferase 1, CD296, ART1

Target/Specificity

This ART1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 14-43 amino acids from the N-terminal region of human ART1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ART1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ART1 Antibody (N-term) - Protein Information**Name** ART1**Function** Has ADP-ribosyltransferase activity toward GLP1R.


Cellular Location

Sarcoplasmic reticulum membrane; Lipid-anchor, GPI-anchor

ART1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ART1 Antibody (N-term) - Images

The anti-ART1 Pab (Cat. #AP2311a) is used in Western blot to detect ART1 in mouse brain tissue lysate.

ART1 Antibody (N-term) - Background

ADP-ribosyltransferase catalyzes the ADP-ribosylation of arginine residues in proteins. Mono-ADP-ribosylation is a posttranslational modification of proteins that is interfered with by a variety of bacterial toxins including cholera, pertussis, and heat-labile enterotoxins of *E. coli*. The amino acid sequence of ART1 consists of predominantly hydrophobic N- and C-terminal regions, which is characteristic of glycosylphosphatidylinositol (GPI)-anchored proteins.

ART1 Antibody (N-term) - References

Koch-Nolte, F., et al., Genomics 39(3):370-376 (1997). Koch-Nolte, F., et al., Genomics 36(1):215-216 (1996). Okazaki, I.J., et al., Biochemistry 33(43):12828-12836 (1994).

ART1 Antibody (N-term) - Citations

- [Regulation of the RhoA/ROCK/AKT/ \$\beta\$ -catenin pathway by arginine-specific ADP-ribosyltransferases 1 promotes migration and epithelial-mesenchymal transition in colon carcinoma.](#)
- [Arginine ADP-ribosyltransferase 1 promotes angiogenesis in colorectal cancer via the](#)

[PI3K/Akt pathway.](#)