

FNTA Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2420a

Specification

FNTA Antibody (Center) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB,E <u>P49354</u> <u>O04631</u>, <u>O61239</u>, <u>P29702</u> Human, Mouse Bovine, Rat Rabbit Polyclonal Rabbit IgG 44409 88-118

FNTA Antibody (Center) - Additional Information

Gene ID 2339

Other Names

Protein farnesyltransferase/geranylgeranyltransferase type-1 subunit alpha, CAAX farnesyltransferase subunit alpha, FTase-alpha, Ras proteins prenyltransferase subunit alpha, Type I protein geranyl-geranyltransferase subunit alpha, GGTase-I-alpha, FNTA

Target/Specificity

This FNTA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-118 amino acids from the Central region of human FNTA.

Dilution WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FNTA Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FNTA Antibody (Center) - Protein Information



Name FNTA

Function Essential subunit of both the farnesyltransferase and the geranylgeranyltransferase complex. Contributes to the transfer of a farnesyl or geranylgeranyl moiety from farnesyl or geranylgeranyl diphosphate to a cysteine at the fourth position from the C-terminus of several proteins having the C-terminal sequence Cys-aliphatic- aliphatic-X. May positively regulate neuromuscular junction development downstream of MUSK via its function in RAC1 prenylation and activation.

FNTA Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

FNTA Antibody (Center) - Images



The anti-FNTA Pab (Cat. #AP2420a) is used in Western blot to detect FNTA in mouse brain tissue lysate.

FNTA Antibody (Center) - Background

FNTA, also known as CAAX farnesyltransferase (FTase), attaches a farnesyl group from farnesyl pyrophosphate to cysteine residues at the fourth position from the C terminus of proteins that end in the so-called CAAX box, where C is cysteine, A is usually but not always an aliphatic amino acid, and X is typically methionine or serine. This type of posttranslational modification provides a mechanism for membrane localization of proteins that lack a transmembrane domain. This enzyme has the remarkable property of farnesylating peptides as short as four residues in length that conform to the CAAX consensus sequence.

FNTA is also a specific cytoplasmic interactor of the transforming growth factor-beta and activin type I receptors. It is likely to be a key component of the signaling pathway which involves p21ras,



an important substrate for farnesyltransferase.

FNTA Antibody (Center) - References

Wang, T., et al., Science 271(5252):1120-1122 (1996). Zhang, F.L., et al., J. Biol. Chem. 269(5):3175-3180 (1994). Andres, D.A., et al., Genomics 18(1):105-112 (1993). Omer, C.A., et al., Biochemistry 32(19):5167-5176 (1993).