

**Prohibitin (PHB1) Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2710a**

**Specification**

---

**Prohibitin (PHB1) Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">P35232</a>
Other Accession	<a href="#">P84173</a> , <a href="#">Q3T165</a>
Reactivity	Human, Mouse
Predicted	Bovine, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit Ig
Calculated MW	29804
Antigen Region	92-121

**Prohibitin (PHB1) Antibody (N-term) - Additional Information**

**Gene ID** 5245

**Other Names**

Prohibitin, PHB

**Target/Specificity**

This Prohibitin (PHB1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 92-121 amino acids from the N-terminal region of human Prohibitin (PHB1).

**Dilution**

WB~~1:2000

IHC-P~~1:100

FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Prohibitin (PHB1) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Prohibitin (PHB1) Antibody (N-term) - Protein Information**

**Name** PHB ([HGNC:8912](#))

**Function** Protein with pleiotropic attributes mediated in a cell- compartment- and tissue-specific manner, which include the plasma membrane-associated cell signaling functions, mitochondrial chaperone, and transcriptional co-regulator of transcription factors in the nucleus (PubMed:[11302691](#), PubMed:[20959514](#), PubMed:[28017329](#), PubMed:[31522117](#)). Plays a role in adipose tissue and glucose Homeostasis in a sex-specific manner (By similarity). Contributes to pulmonary vascular remodeling by accelerating proliferation of pulmonary arterial smooth muscle cells (By similarity).

**Cellular Location**

Mitochondrion inner membrane. Nucleus. Cytoplasm. Cell membrane

**Tissue Location**


Widely expressed in different tissues.

**Prohibitin (PHB1) Antibody (N-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Prohibitin (PHB1) Antibody (N-term) - Images**




All lanes : Anti-Prohibitin (PHB1) Antibody (N-term) at 1:2000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: Jurkat whole cell lysate Lane 5: HeLa whole cell lysate Lane 6: Mouse kidney tissue lysate Lane 7: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 30 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of AP2710A on paraffin-embedded Human liver tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP2710A on paraffin-embedded Human kidney tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Flow cytometric analysis of ATDC5 cells using PHB1 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **Prohibitin (PHB1) Antibody (N-term) - Background**

Prohibitin is an evolutionarily conserved protein that is ubiquitously expressed. It is thought to be a negative regulator of cell proliferation and may be a tumor suppressor. Mutations have been linked to sporadic breast cancer. Prohibitin is expressed as two transcripts with varying lengths of 3' untranslated region.

### **Prohibitin (PHB1) Antibody (N-term) - References**

Gregory-Bass,R.C., Int. J. Cancer 122 (9), 1923-1930 (2008) Ross,J.A., J. Biol. Chem. 283 (8), 4699-4713 (2008) White,J.J., Genomics 11 (1), 228-230 (1991)