

M Sema5a Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2712a

Specification

M Sema5a Antibody (N-term) - Product Information

Application WB,E
Primary Accession O13591

Other Accession <u>D3ZTD8</u>, <u>Q62217</u>

Reactivity
Predicted
Host
Clonality
Isotype

Mouse
Rat
Rabbit
Polyclonal
Rabbit IgG

M Sema5a Antibody (N-term) - Additional Information

Gene ID 9037

Other Names

Semaphorin-5A, Semaphorin-F, Sema F, SEMA5A, SEMAF

Target/Specificity

This Mouse Sema5a antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of mouse Sema5a.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

M Sema5a Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

M Sema5a Antibody (N-term) - Protein Information

Name SEMA5A

Synonyms SEMAF



Function Bifunctional axonal guidance cue regulated by sulfated proteoglycans; attractive effects result from interactions with heparan sulfate proteoglycans (HSPGs), while the inhibitory effects depend on interactions with chondroitin sulfate proteoglycans (CSPGs) (By similarity). Ligand for receptor PLXNB3. In glioma cells, SEMA5A stimulation of PLXNB3 results in the disassembly of F-actin stress fibers, disruption of focal adhesions and cellular collapse as well as inhibition of cell migration and invasion through ARHGDIA-mediated inactivation of RAC1. May promote angiogenesis by increasing endothelial cell proliferation and migration and inhibiting apoptosis.

Cellular Location


Membrane; Single-pass type I membrane protein.

M Sema5a Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

M Sema5a Antibody (N-term) - Images




Anti-M Sema5a Antibody (N-term) at 1:1000 dilution + Mouse liver tissue lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 121 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

M Sema5a Antibody (N-term) - Background

Members of the semaphorin protein family, such as SEMA5A, are involved in axonal guidance during neural development.

M Sema5a Antibody (N-term) - References





Tel: 858.875.1900 Fax: 858.875.1999

Bialecka, M., Neurosci. Lett. 399 (1-2), 121-123 (2006) Melin, M., Neuropsychobiology 54 (1), 64-69 (2006) M Sema5a Antibody (N-term) - Citations

• Identification of a novel biomarker, SEMA5A, for non-small cell lung carcinoma in nonsmoking women.