

## **INDO Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2728c

# **Specification**

## **INDO Antibody (Center) - Product Information**

**Application** IHC-P, WB,E **Primary Accession** P14902 Other Accession NP 002155.1 **Human**, Mouse Reactivity Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 45326 Antigen Region 79-105

## INDO Antibody (Center) - Additional Information

#### **Gene ID 3620**

### **Other Names**

Indoleamine 2, 3-dioxygenase 1, IDO-1, Indoleamine-pyrrole 2, 3-dioxygenase, IDO1, IDO, INDO

### Target/Specificity

This INDO antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 79-105 amino acids from the Central region of human INDO.

### **Dilution**

IHC-P~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

INDO Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## **INDO Antibody (Center) - Protein Information**

Name IDO1 (<u>HGNC:6059</u>)



# Synonyms IDO, INDO

**Function** Catalyzes the first and rate limiting step of the catabolism of the essential amino acid tryptophan along the kynurenine pathway (PubMed:<u>17671174</u>). Involved in the peripheral immune tolerance, contributing to maintain homeostasis by preventing autoimmunity or immunopathology that would result from uncontrolled and overreacting immune responses (PubMed:<u>25691885</u>). Tryptophan shortage inhibits T lymphocytes division and accumulation of tryptophan catabolites induces T-cell apoptosis and differentiation of regulatory T-cells (PubMed:<u>25691885</u>). Acts as a suppressor of anti-tumor immunity (PubMed:<u>14502282</u>, PubMed:<u>23103127</u>, PubMed:<u>25157255</u>, PubMed:<u>25691885</u>). Limits the growth of intracellular pathogens by depriving tryptophan (PubMed:<u>25691885</u>). Protects the fetus from maternal immune rejection (PubMed:<u>25691885</u>).

#### **Cellular Location**

Cytoplasm, cytosol {ECO:0000250|UniProtKB:P28776, ECO:0000303|PubMed:25691885}

### **Tissue Location**

Expressed in mature dendritic cells located in lymphoid organs (including lymph nodes, spleen, tonsils, Peyers's patches, the gut lamina propria, and the thymic medulla), in some epithelial cells of the female genital tract, as well as in endothelial cells of term placenta and in lung parenchyma (PubMed:25691885). Weakly or not expressed in most normal tissues, but mostly inducible in most tissues (PubMed:25691885). Expressed in more than 50% of tumors, either by tumoral, stromal, or endothelial cells (expression in tumor is associated with a worse clinical outcome) (PubMed:18418598). Not overexpressed in tumor-draining lymph nodes (PubMed:25691885, PubMed:26155395).


# **INDO Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## INDO Antibody (Center) - Images







INDO Antibody (Center) (Cat. #AP2728c) western blot analysis in mouse cerebellum tissue lysates (35ug/lane). This demonstrates the INDO antibody detected the INDO protein (arrow).



INDO Antibody (Center) (Cat. #AP2728c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the INDO antibody detected the INDO protein (arrow).







Tel: 858.875.1900 Fax: 858.875.1999

INDO antibody (Center) (Cat. #AP2728c) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the INDO antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.