

**ITGA6 Antibody (isoform 2 S1064)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2753A****Specification**

---

**ITGA6 Antibody (isoform 2 S1064) - Product Information**

Application	IF, WB, IHC-P,E
Primary Accession	<a href="#">P23229-2</a>
Other Accession	<a href="#">P26007</a>
Reactivity	Human
Predicted	Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1043-1072

**ITGA6 Antibody (isoform 2 S1064) - Additional Information****Other Names**

Integrin alpha-6, CD49 antigen-like family member F, VLA-6, CD49f, Integrin alpha-6 heavy chain, Integrin alpha-6 light chain, Processed integrin alpha-6, Alpha6p, ITGA6

**Target/Specificity**

This ITGA6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1043-1072 amino acids from isoform 2 of human ITGA6.

**Dilution**

IF~~1:200  
WB~~1:1000  
IHC-P~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**


ITGA6 Antibody (isoform 2 S1064) is for research use only and not for use in diagnostic or therapeutic procedures.

**ITGA6 Antibody (isoform 2 S1064) - Protein Information****ITGA6 Antibody (isoform 2 S1064) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### ITGA6 Antibody (isoform 2 S1064) - Images




Fluorescent confocal image of HeLa cells stained with ITGA6 (isoform 2 S1064) antibody. HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP2753a ITGA6 (isoform 2 S1064) primary antibody (1:200, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). Note the highly specific localization of the ITGA6 mainly to the nucleus, supported by Human Protein Atlas Data (<http://www.proteinatlas.org/ENSG00000091409>).



Anti-ITGA6 Antibody (isoform 2 S1064) at 1:1000 dilution + SW480 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 119 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot analysis of anti-ITGA6 Antibody (isoform 2 S1064) (Cat.#AP2753a) in 293 cell line lysates (35ug/lane). ITGA6 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with ITGA6 Antibody (isoform 2 S1064) (Cat.#AP2753a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

#### **ITGA6 Antibody (isoform 2 S1064) - Background**

The ITGA6 protein product is the integrin alpha chain alpha 6. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. A given chain may combine with multiple partners resulting in different integrins. For example, alpha 6 may combine with beta 4 in the integrin referred to as TSP180, or with beta 1 in the integrin VLA-6. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling.

#### **ITGA6 Antibody (isoform 2 S1064) - References**

References for protein:

1. Yang, X.H., Cancer Res. 68 (9), 3204-3213 (2008)
2. Hayashi, R., Biochem. Biophys. Res. Commun. 367 (2), 256-263 (2008)

References for HeLa cell line:

1. Scherer WF, Syverton JT, Gey GO (May 1953). "Studies on the propagation in vitro of poliomyelitis viruses. IV. Viral multiplication in a stable strain of human malignant epithelial cells (strain HeLa) derived from an epidermoid carcinoma of the cervix". J. Exp. Med. 97 (5): 695-710. [PubMed:13052828].
2. Macville M, Schröck E, Padilla-Nash H, Keck C, Ghadimi BM, Zimonjic D, Popescu N, Ried T (January 1999). "Comprehensive and definitive molecular cytogenetic characterization of HeLa cells by spectral karyotyping". Cancer Res. 59 (1): 141-50. [PubMed: 9892199].
3. Rahbari R, Sheahan T, Modes V, Collier P, Macfarlane C, Badge RM (April 2009). "A novel L1 retrotransposon marker for HeLa cell line identification". BioTechniques 46 (4): 277-84. [PubMed: 19450234].
4. Capes-Davis A, Theodosopoulos G, Atkin I, Drexler HG, Kohara A, MacLeod RA, Masters JR, Nakamura Y, Reid YA, Reddel RR, Freshney RI (July 2010). "Check your cultures! A list of cross-contaminated or misidentified cell lines". Int. J. Cancer 127 (1): 1-8. [PubMed:20143388].