

**B2M Antibody (C-term Y86)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2771d**

### Specification

#### B2M Antibody (C-term Y86) - Product Information

Application	WB,E
Primary Accession	<a href="#">P61769</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	13715
Antigen Region	71-98

#### B2M Antibody (C-term Y86) - Additional Information

##### Gene ID 567

##### Other Names

Beta-2-microglobulin, Beta-2-microglobulin form pl 53, B2M

##### Target/Specificity

This B2M antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 71-98 amino acids from the C-terminal region of human B2M.

##### Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

B2M Antibody (C-term Y86) is for research use only and not for use in diagnostic or therapeutic procedures.

#### B2M Antibody (C-term Y86) - Protein Information

##### Name B2M ([HGNC:914](#))

**Function** Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system. Exogenously applied M.tuberculosis EsxA

or EsxA-EsxB (or EsxA expressed in host) binds B2M and decreases its export to the cell surface (total protein levels do not change), probably leading to defects in class I antigen presentation (PubMed:[25356553](#)).

#### Cellular Location


Secreted. Cell surface. Note=Detected in serum and urine (PubMed:1336137, PubMed:7554280). {ECO:0000269|PubMed:7554280, ECO:0000269|Ref.6}

#### B2M Antibody (C-term Y86) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### B2M Antibody (C-term Y86) - Images



B2M Antibody (C-term) (Cat. #AP2771d) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the B2M antibody detected the B2M protein (arrow).

#### B2M Antibody (C-term Y86) - Background

Beta-2-microglobulin is a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells.

#### B2M Antibody (C-term Y86) - References

Sakata,M., J. Mol. Biol. 382 (5), 1242-1255 (2008)  
Huang,W.C., Clin. Cancer Res. 14 (17), 5341-5347 (2008)  
Gattoni-Celli,S., Cancer Res. 52 (5), 1201-1204 (1992)