

HNF4A Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2778c

Specification

HNF4A Antibody (Center) - Product Information

Application WB, FC, IF, IHC-P,E

Primary Accession P41235

Other Accession
Reactivity
Reactivity
Predicted
Host
Clonality
Isotype
Rabbit IgG

Calculated MW 52785
Antigen Region 281-312

HNF4A Antibody (Center) - Additional Information

Gene ID 3172

Other Names

Hepatocyte nuclear factor 4-alpha, HNF-4-alpha, Nuclear receptor subfamily 2 group A member 1, Transcription factor 14, TCF-14, Transcription factor HNF-4, HNF-4, HNF-4, HNF-4, TCF-14

Target/Specificity

This HNF4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 281-312 amino acids from the Central region of human HNF4A.

Dilution

WB~~1:1000 FC~~1:10~50 IF~~1:10~50 IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HNF4A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HNF4A Antibody (Center) - Protein Information



Name HNF4A

Synonyms HNF4, NR2A1, TCF14

Function Transcriptional regulator which controls the expression of hepatic genes during the transition of endodermal cells to hepatic progenitor cells, facilitating the recruitment of RNA pol II to the promoters of target genes (PubMed:30597922). Activates the transcription of CYP2C38 (By similarity). Represses the CLOCK-BMAL1 transcriptional activity and is essential for circadian rhythm maintenance and period regulation in the liver and colon cells (PubMed:30530698).


Cellular Location Nucleus.

HNF4A Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

HNF4A Antibody (Center) - Images




Confocal immunofluorescent analysis of HNF4A Antibody (Center)(Cat#AP2778c) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red).DAPI was used to stain the cell nuclear (blue).






HNF4A Antibody (Center) (Cat. #AP2778c) western blot analysis in 293,HepG2 cell line and rat liver tissue lysates (35ug/lane).This demonstrates the HNF4A antibody detected the HNF4A protein (arrow).



HNF4A Antibody (Center) (AP2778c)immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of HNF4A Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.





HNF4A Antibody (Center) (Cat. #AP2778c) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HNF4A Antibody (Center) - Background

HNF4A is a nuclear transcription factor which binds DNA as a homodimer. This protein controls the expression of several genes, including hepatocyte nuclear factor 1 alpha, a transcription factor which regulates the expression of several hepatic genes. This protein may play a role in development of the liver, kidney, and intestines. Mutations in HNF4A gene have been associated with monogenic autosomal dominant non-insulin-dependent diabetes mellitus type I.

HNF4A Antibody (Center) - References

Kritis A.A., Argyrokastritis A.Gene 173:275-280(1996) Yamagata K., Furuta H., Oda N.Nature 384:458-460(1996) Furuta H., Iwasaki N., Oda N.Diabetes 46:1652-1657(1997) Moeller A.M., Urhammer S.A.Diabetologia 40:980-983(1997)