

SFRS2 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2800c**Specification**

SFRS2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q01130
Other Accession	Q6PDU1 , Q06A98 , Q62093 , P30352 , Q3MHR5
Reactivity	Human
Predicted	Bovine, Chicken, Mouse, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25476
Antigen Region	76-105

SFRS2 Antibody (Center) - Additional Information**Gene ID** 6427**Other Names**

Serine/arginine-rich splicing factor 2, Protein PR264, Splicing component, 35 kDa, Splicing factor SC35, SC-35, Splicing factor, arginine/serine-rich 2, SRSF2, SFRS2

Target/Specificity

This SFRS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 76-105 amino acids from the Central region of human SFRS2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SFRS2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SFRS2 Antibody (Center) - Protein Information**Name** SRSF2

Synonyms SFRS2

Function Necessary for the splicing of pre-mRNA. It is required for formation of the earliest ATP-dependent splicing complex and interacts with spliceosomal components bound to both the 5'- and 3'-splice sites during spliceosome assembly. It also is required for ATP-dependent interactions of both U1 and U2 snRNPs with pre-mRNA. Interacts with other spliceosomal components, via the RS domains, to form a bridge between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF. Binds to purine-rich RNA sequences, either 5'-AGSAGAGTA-3' (S=C or G) or 5'-GTTCGAGTA-3'. Can bind to beta-globin mRNA and commit it to the splicing pathway. The phosphorylated form (by SRPK2) is required for cellular apoptosis in response to cisplatin treatment.

Cellular Location


Nucleus. Nucleus, nucleoplasm. Nucleus speckle. Note=Phosphorylation by SRPK2 provokes its redistribution from the nuclear speckle to nucleoplasm

SFRS2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SFRS2 Antibody (Center) - Images



Western blot analysis of SFRS2 Antibody (Center) (Cat. #AP2800c) in mouse stomach tissue lysates (35ug/lane). SFRS2 (arrow) was detected using the purified Pab.

SFRS2 Antibody (Center) - Background

SFRS2 is necessary for the splicing of pre-mRNA. The protein is required for formation of the earliest ATP-dependent splicing complex and interacts with spliceosomal components bound to both the 5'- and 3'-splice sites during spliceosome assembly. It also is required for ATP-dependent interactions of both U1 and U2 snRNPs with pre-mRNA. And it interacts with other spliceosomal

components, via the RS domains, to form a bridge between the 5'- and 3'-splice site binding components, U1 snRNP and U2AF. It binds to purine-rich RNA sequences, either 5'-AGSAGAGTA-3' (S=C or G) or 5'-GTTCGAGTA-3' and can bind to beta-globin mRNA and commit it to the splicing pathway.

SFRS2 Antibody (Center) - References

Merdzhanova,G., Cell Death Differ. 15 (12), 1815-1823 (2008)
Solis,A.S., J. Biol. Chem. 283 (35), 23619-23626 (2008)
Donev,R., Mol. Psychiatry 12 (7), 681-690 (2007)
Sureau,A., Proc. Natl. Acad. Sci. U.S.A. 89 (24), 11683-11687 (1992)