

**HSPH1 Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2860d****Specification**

---

**HSPH1 Antibody (Center) - Product Information**

Application	FC, WB, IHC-P,E
Primary Accession	<a href="#">Q92598</a>
Other Accession	<a href="#">Q66HA8</a> , <a href="#">Q60446</a> , <a href="#">Q0IIM3</a>
Reactivity	Human
Predicted	Bovine, Hamster, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	96865
Antigen Region	549-579

**HSPH1 Antibody (Center) - Additional Information****Gene ID** 10808**Other Names**

Heat shock protein 105 kDa, Antigen NY-CO-25, Heat shock 110 kDa protein, HSPH1, HSP105, HSP110, KIAA0201

**Target/Specificity**

This HSPH1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 549-579 amino acids from the Central region of human HSPH1.

**Dilution**

FC~~1:10~50

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

HSPH1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**HSPH1 Antibody (Center) - Protein Information**

**Name** HSPH1

**Synonyms** HSP105, HSP110, KIAA0201

**Function** Acts as a nucleotide-exchange factor (NEF) for chaperone proteins HSPA1A and HSPA1B, promoting the release of ADP from HSPA1A/B thereby triggering client/substrate protein release (PubMed:[24318877](#)). Prevents the aggregation of denatured proteins in cells under severe stress, on which the ATP levels decrease markedly. Inhibits HSPA8/HSC70 ATPase and chaperone activities (By similarity).

**Cellular Location**

Cytoplasm.

**Tissue Location**


Highly expressed in testis. Present at lower levels in most brain regions, except cerebellum. Overexpressed in cancer cells.

**HSPH1 Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**HSPH1 Antibody (Center) - Images**



Western blot analysis of HSPH1 Antibody (Center) (Cat. #AP2860d) in K562 cell line lysates (35ug/lane). HSPH1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human testis tissue reacted with HSPH1 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



HSPH1 Antibody (Center) (Cat. #AP2860d) flow cytometric analysis of k562 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **HSPH1 Antibody (Center) - Background**

HSPH1 prevents the aggregation of denatured proteins in cells under severe stress, on which the ATP levels decrease markedly. This protein inhibits HSPA8/HSC70 ATPase and chaperone activities.

### **HSPH1 Antibody (Center) - References**

- Ishihara K., Yasuda K., Hatayama T. Biochim. Biophys. Acta 1444:138-142(1999)
- Nagase T., Seki N., Ishikawa K., Ohira M., Kawarabayashi Y., DNA Res. 3:321-329(1996)
- The MGC Project Team Genome Res. 14:2121-2127(2004)
- Miyazaki M., Nakatsura T., Yokomine K., Senju S., Monji M., Hosaka S., Cancer Sci. 96:695-705(2005)