

**ELP2 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP2884b****Specification**

---

**ELP2 Antibody (C-term) - Product Information**

Application	FC, WB, IF,E
Primary Accession	<a href="#">Q6IA86</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	92500
Antigen Region	737-765

**ELP2 Antibody (C-term) - Additional Information****Gene ID** 55250**Other Names**

Elongator complex protein 2, ELP2, SHINC-2, STAT3-interacting protein 1, StIP1, ELP2, STATIP1

**Target/Specificity**

This ELP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 737-765 amino acids from the C-terminal region of human ELP2.

**Dilution**

FC~~1:10~50

WB~~1:1000

IF~~1:100

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ELP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ELP2 Antibody (C-term) - Protein Information****Name** ELP2

## Synonyms STATIP1

**Function** Component of the elongator complex which is required for multiple tRNA modifications, including mcm5U (5-methoxycarbonylmethyl uridine), mcm5s2U (5-methoxycarbonylmethyl-2-thiouridine), and ncm5U (5-carbamoylmethyl uridine) (PubMed:[29332244](#)). The elongator complex catalyzes the formation of carboxymethyluridine in the wobble base at position 34 in tRNAs (PubMed:[29332244](#)).

## Cellular Location


Cytoplasm. Nucleus

## ELP2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## ELP2 Antibody (C-term) - Images



Fluorescent confocal image of SY5Y cells stained with ELP2 (C-term) antibody. SY5Y cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min). Cells were then incubated with AP2884b ELP2 (C-term) primary antibody (1:100, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 5 min). Note the highly specific localization of the ELP2 immunosignal mainly to the cytoplasm, supported by Human Protein Atlas Data (<http://www.proteinatlas.org/ENSG00000134759>).



Western blot analysis of ELP2 antibody (C-term) (Cat. #AP2884b) in NCI-H460 cell line lysates (35ug/lane). ELP2 (arrow) was detected using the purified Pab.



Flow cytometric analysis of NCI-H292 cells using ELP2 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### ELP2 Antibody (C-term) - Background

ELP2 regulates the ligand-dependent activation of STAT3. The protein acts as subunit of the RNA polymerase II elongator complex, which is a histone acetyltransferase component of the RNA polymerase II (Pol II) holoenzyme and is involved in transcriptional elongation. Elongator may play a role in chromatin remodeling and is involved in acetylation of histones H3 and probably H4.

### ELP2 Antibody (C-term) - References

References for protein:

1.Kim,J.H., Proc. Natl. Acad. Sci. U.S.A. 99 (3), 1241-1246 (2002)

2.Hawkes,N.A., J. Biol. Chem. 277 (4), 3047-3052 (2002)

References for SY5Y (SH-SY5Y; ATCC#CRL-2266): 1. Ross RA, et al. Coordinate morphological and biochemical interconversion of human neuroblastoma cells. J. Natl. Cancer Inst. 71: 741-749, 1983. [PubMed: 6137586]; 2. Biedler JL, et al. Multiple neurotransmitter synthesis by human neuroblastoma cell lines and clones. Cancer Res. 38: 3751-3757, 1978. [PubMed: 29704].