

CCT3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2890b**Specification**

CCT3 Antibody (C-term) - Product Information

Application	FC, WB, IHC-P,E
Primary Accession	P49368
Other Accession	Q4R963 , Q3T0K2
Reactivity	Human
Predicted	Bovine, Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	60534
Antigen Region	509-536

CCT3 Antibody (C-term) - Additional Information**Gene ID** 7203**Other Names**

T-complex protein 1 subunit gamma, TCP-1-gamma, CCT-gamma, hTRiC5, CCT3, CCTG, TRiC5

Target/Specificity

This CCT3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 509-536 amino acids from the C-terminal region of human CCT3.

Dilution

FC~~1:10~50

WB~~1:1000

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCT3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CCT3 Antibody (C-term) - Protein Information

Name CCT3

Synonyms CCTG, TRIC5

Function Component of the chaperonin-containing T-complex (TRiC), a molecular chaperone complex that assists the folding of actin, tubulin and other proteins upon ATP hydrolysis (PubMed:[25467444](#), PubMed:[36493755](#), PubMed:[35449234](#), PubMed:[37193829](#)). The TRiC complex mediates the folding of WRAP53/TCAB1, thereby regulating telomere maintenance (PubMed:[25467444](#)). As part of the TRiC complex may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia (PubMed:[20080638](#)).

Cellular Location


Cytoplasm.

CCT3 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CCT3 Antibody (C-term) - Images



Western blot analysis of CCT3 Antibody (C-term) (Cat. #AP2890b) in K562 cell line lysates (35ug/lane). CCT3 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with CCT3 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of K562 cells using CCT3 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram) FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CCT3 Antibody (C-term) - Background

CCT3 is a molecular chaperone that is member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of two identical stacked rings, each containing eight different proteins. Unfolded polypeptides enter the central cavity of the complex and are folded in an ATP-dependent manner. The complex folds various proteins, including actin and tubulin.

CCT3 Antibody (C-term) - References

Zebol,J.R., Int. J. Biochem. Cell Biol. 41 (4), 822-827 (2009)