

**Phospho-Bid(S65) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3041a**

## Specification

---

### Phospho-Bid(S65) Antibody - Product Information

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P55957</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>21995</b>

### Phospho-Bid(S65) Antibody - Additional Information

**Gene ID** 637

#### Other Names

BH3-interacting domain death agonist, p22 BID, BID, BH3-interacting domain death agonist p15, p15 BID, BH3-interacting domain death agonist p13, p13 BID, BH3-interacting domain death agonist p11, p11 BID, BID

#### Target/Specificity

This Bid Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S65 of human Bid.

#### Dilution

WB~~1:1000  
IHC-P~~1:50~100  
E~~Use at an assay dependent concentration.

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Phospho-Bid(S65) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Phospho-Bid(S65) Antibody - Protein Information

**Name** BID

**Function** Induces caspases and apoptosis (PubMed:[14583606](#)). Counters the protective effect of BCL2 (By similarity).

#### Cellular Location

Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane. Note=When uncleaved, it is predominantly cytoplasmic. [BH3-interacting domain death agonist p13]: Mitochondrion membrane {ECO:0000250|UniProtKB:P70444}. Note=Associated with the mitochondrial membrane. {ECO:0000250|UniProtKB:P70444} [Isoform 3]: Cytoplasm

#### Tissue Location


[Isoform 2]: Expressed in spleen, pancreas and placenta (at protein level). [Isoform 4]: Expressed in lung and pancreas (at protein level).

### Phospho-Bid(S65) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Phospho-Bid(S65) Antibody - Images



The anti-Phospho-Bid-S65 Pab (Cat. #AP3041a) is used in Western blot to detect Phospho-Bid-S65 in Jurkat tissue lysate



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

#### **Phospho-Bid(S65) Antibody - Background**

Bid is a death agonist that heterodimerizes with either agonist BAX or antagonist BCL2. The encoded protein is a member of the BCL-2 family of cell death regulators. Bid induces ICE-like proteases and apoptosis. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the major proteolytic product p15 Bid translocates to mitochondria where it triggers cytochrome c release.

#### **Phospho-Bid(S65) Antibody - References**

Liu, J., et al., *Biochem. Biophys. Res. Commun.* 330(3):865-870 (2005).  
Broaddus, V.C., et al., *J. Biol. Chem.* 280(13):12486-12493 (2005).  
Weng, C., et al., *J. Biol. Chem.* 280(11):10491-10500 (2005).  
Gong, X.M., et al., *J. Biol. Chem.* 279(28):28954-28960 (2004).  
Garcia-Saez, A.J., et al., *Biochemistry* 43(34):10930-10943 (2004).