

**Phospho-GATA6(Y271) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3536a****Specification**

---

**Phospho-GATA6(Y271) Antibody - Product Information**

Application	WB, DB, E
Primary Accession	<a href="#">O92908</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

**Phospho-GATA6(Y271) Antibody - Additional Information****Gene ID** 2627**Other Names**

Transcription factor GATA-6, GATA-binding factor 6, GATA6

**Target/Specificity**

This GATA6 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y271 of human GATA6.

**Dilution**

WB~~1:1000

DB~~1:500

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Phospho-GATA6(Y271) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-GATA6(Y271) Antibody - Protein Information****Name** GATA6

**Function** Transcriptional activator (PubMed:[19666519](#), PubMed:[22750565](#), PubMed:[22824924](#), PubMed:[27756709](#)). Regulates SEMA3C and PLXNA2 (PubMed:[19666519](#)). Involved in gene regulation specifically in the gastric epithelium (PubMed:[9315713](#)). May regulate genes that

protect epithelial cells from bacterial infection (PubMed:[16968778](#)). Involved in bone morphogenetic protein (BMP)-mediated cardiac-specific gene expression (By similarity). Binds to BMP response element (BMPRE) DNA sequences within cardiac activating regions (By similarity). In human skin, controls several physiological processes contributing to homeostasis of the upper pilosebaceous unit. Triggers ductal and sebaceous differentiation as well as limits cell proliferation and lipid production to prevent hyperseborrhoea. Mediates the effects of retinoic acid on sebocyte proliferation, differentiation and lipid production. Also contributes to immune regulation of sebocytes and antimicrobial responses by modulating the expression of anti-inflammatory genes such as IL10 and pro-inflammatory genes such as IL6, TLR2, TLR4, and IFNG. Activates TGFB1 signaling which controls the interfollicular epidermis fate (PubMed:[33082341](#)).

#### **Cellular Location**

Nucleus

#### **Tissue Location**


Expressed in heart, gut and gut-derived tissues. Expressed in skin upper pilosebaceous unit. Expression is decreased or lost in acne lesions (PubMed:33082341).

### **Phospho-GATA6(Y271) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Phospho-GATA6(Y271) Antibody - Images**



Dot blot analysis of anti-Phospho-GATA6-pY271 Phospho-specific Pab (RB15308) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

### **Phospho-GATA6(Y271) Antibody - Background**

GATA6 may be important for regulating terminal differentiation and/or proliferation.

**Phospho-GATA6(Y271) Antibody - References**

Kwei,K.A., PLoS Genet. 4 (5), E1000081 (2008)  
Ghatnekar,A., Biochim. Biophys. Acta 1779 (3), 145-151 (2008)