

# Phospho-PBK(Y74) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3577a

## **Specification**

# Phospho-PBK(Y74) Antibody - Product Information

Application DB,E
Primary Accession Q96KB5
Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG

# Phospho-PBK(Y74) Antibody - Additional Information

### Gene ID 55872

#### **Other Names**

Lymphokine-activated killer T-cell-originated protein kinase, Cancer/testis antigen 84, CT84, MAPKK-like protein kinase, Nori-3, PDZ-binding kinase, Spermatogenesis-related protein kinase, SPK, T-LAK cell-originated protein kinase, PBK, TOPK

# **Target/Specificity**

This PBK Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Y74 of human PBK.

#### **Dilution**

DB~~1:500

E~~Use at an assay dependent concentration.

## **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

## Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

Phospho-PBK(Y74) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# Phospho-PBK(Y74) Antibody - Protein Information

# **Name PBK**

## **Synonyms** TOPK



Function Phosphorylates MAP kinase p38. Seems to be active only in mitosis. May also play a role in the activation of lymphoid cells. When phosphorylated, forms a complex with TP53, leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin- induced DNA damage.

### **Tissue Location**


Expressed in the testis and placenta. In the testis, restrictedly expressed in outer cell layer of seminiferous tubules.

## Phospho-PBK(Y74) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Phospho-PBK(Y74) Antibody - Images



Dot blot analysis of anti-Phospho-PBK-Y74 Antibody (Cat.#AP3577a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

# Phospho-PBK(Y74) Antibody - Background

PBK is a serine/threonine kinase related to the dual specific mitogen-activated protein kinase kinase (MAPKK) family. Evidence suggests that mitotic phosphorylation is required for its catalytic activity. This mitotic kinase may be involved in the activation of lymphoid cells and support testicular functions, with a suggested role in the process of spermatogenesis.

## Phospho-PBK(Y74) Antibody - References





Tel: 858.875.1900 Fax: 858.875.1999

Nandi, A., et al., Blood Cells Mol. Dis. 32(1):240-245 (2004). Simons-Evelyn, M., et al., Blood Cells Mol. Dis. 27(5):825-829 (2001). Zhao, S., et al., Int. J. Biochem. Cell Biol. 33(6):631-636 (2001). Abe, Y., et al., J. Biol. Chem. 275(28):21525-21531 (2000). Gaudet, S., et al., Proc. Natl. Acad. Sci. U.S.A. 97(10):5167-5172 (2000).

# Phospho-PBK(Y74) Antibody - Citations

- TOPK/PBK is phosphorylated by ERK2 at serine 32, promotes tumorigenesis and is involved in sorafenib resistance in RCC
- Phosphorylation of TOPK at Y74, Y272 by Src increases the stability of TOPK and promotes tumorigenesis of colon