

**Phospho-CDX2(S283) Antibody**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP3701a**

**Specification**

---

**Phospho-CDX2(S283) Antibody - Product Information**

Application	WB, DB, IF, E
Primary Accession	<a href="#">Q99626</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

**Phospho-CDX2(S283) Antibody - Additional Information**

**Gene ID** 1045

**Other Names**

Homeobox protein CDX-2, CDX-3, Caudal-type homeobox protein 2, CDX2, CDX3

**Target/Specificity**

This CDX2 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S283 of human CDX2.

**Dilution**

WB~~1:1000

DB~~1:500

IF~~1:10~50

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Phospho-CDX2(S283) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Phospho-CDX2(S283) Antibody - Protein Information**

**Name** CDX2

**Synonyms** CDX3

**Function** Transcription factor which regulates the transcription of multiple genes expressed in the intestinal epithelium (By similarity). Binds to the promoter of the intestinal sucrase-isomaltase SI and activates SI transcription (By similarity). Binds to the DNA sequence 5'-ATAAAACTTAT-3' in the promoter region of VDR and activates VDR transcription (By similarity). Binds to and activates transcription of LPH (By similarity). Activates VDR transcription of CLDN2 and intestinal mucin MUC2 (By similarity). Binds to the 5'-AATTTTTTACAACACCT-3' DNA sequence in the promoter region of CA1 and activates CA1 transcription (By similarity). Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed:[28473536](#)).

**Cellular Location**

Nucleus {ECO:0000250|UniProtKB:P43241}.

**Tissue Location**


Detected in small intestine, colon and pancreas.

**Phospho-CDX2(S283) Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Phospho-CDX2(S283) Antibody - Images**



Confocal immunofluorescent analysis of Phospho-CDX2-S283 Antibody (Cat#AP3701a) with WiDr cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red).



Dot blot analysis of anti-Phospho-CDX2 Phospho-specific Pab (Cat. #AP3701a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

#### **Phospho-CDX2(S283) Antibody - Background**

The level and beta-cell specificity of insulin gene expression are regulated by a set of nuclear proteins that bind to specific sequences within the promoter of the insulin gene (INS; MIM 176730) and interact with RNA polymerase to activate or repress transcription. The proteins LMX1 (MIM 600298) and CDX3 are homeodomain proteins that bind an A/T-rich sequence in the insulin promoter and stimulate its transcription.

#### **Phospho-CDX2(S283) Antibody - References**

Benoit, Y.D., et al. Am. J. Physiol. Gastrointest. Liver Physiol. 298 (4), G504-G517 (2010)  
Xie, Y., et al. Int. J. Oncol. 36(2):509-516(2010)  
Park do, Y., et al. Mod. Pathol. 23(1):54-61(2010)  
Lora, V., et al. Anticancer Res. 29(12):5033-5037(2009)  
Porjazova, E., et al. Akush Ginekol (Sofia) 48(4):32-34(2009)