

Phospho-FOXP3(S418) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3731a

Specification

Phospho-FOXP3(S418) Antibody - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype DB,E <u>O9BZS1</u> <u>O6U8D7</u>, <u>NP_054728</u> Human Monkey Rabbit Polyclonal Rabbit IgG

Phospho-FOXP3(S418) Antibody - Additional Information

Gene ID 50943

Other Names Forkhead box protein P3, Scurfin, FOXP3, IPEX

Target/Specificity

This FOXP3 Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S418 of human FOXP3.

Dilution DB~~1:500 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Phospho-FOXP3(S418) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-FOXP3(S418) Antibody - Protein Information

Name FOXP3

Synonyms IPEX



Function Transcriptional regulator which is crucial for the development and inhibitory function of regulatory T-cells (Treg) (PubMed:17377532, PubMed:21458306, PubMed:23947341, PubMed:24354325, PubMed:24722479, PubMed:24835996, PubMed:30513302, PubMed:32644293). Plays an essential role in maintaining homeostasis of the immune system by allowing the acquisition of full suppressive function and stability of the Treg lineage, and by directly modulating the expansion and function of conventional T-cells (PubMed: 23169781). Can act either as a transcriptional repressor or a transcriptional activator depending on its interactions with other transcription factors, histone acetylases and deacetylases (PubMed: 17377532, PubMed:21458306, PubMed:23947341, PubMed:24354325, PubMed:24722479). The suppressive activity of Treg involves the coordinate activation of many genes, including CTLA4 and TNFRSF18 by FOXP3 along with repression of genes encoding cytokines such as interleukin-2 (IL2) and interferon-gamma (IFNG) (PubMed:17377532, PubMed:21458306, PubMed:23947341, PubMed:24354325, PubMed:24722479). Inhibits cytokine production and T-cell effector function by repressing the activity of two key transcription factors, RELA and NFATC2 (PubMed: 15790681). Mediates transcriptional repression of IL2 via its association with histone acetylase KAT5 and histone deacetylase HDAC7 (PubMed: 17360565). Can activate the expression of TNFRSF18, IL2RA and CTLA4 and repress the expression of IL2 and IFNG via its association with transcription factor RUNX1 (PubMed:<u>17377532</u>). Inhibits the differentiation of IL17 producing helper T-cells (Th17) by antagonizing RORC function, leading to down-regulation of IL17 expression, favoring Treg development (PubMed: 18368049). Inhibits the transcriptional activator activity of RORA (PubMed:<u>18354202</u>). Can repress the expression of IL2 and IFNG via its association with transcription factor IKZF4 (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:17360565, ECO:0000269|PubMed:18354202, ECO:0000269|PubMed:22678915, ECO:0000269|PubMed:23396208, ECO:0000269|PubMed:23973222, ECO:0000269|PubMed:23973223, ECO:0000269|PubMed:32644293}. Cytoplasm Note=Predominantly expressed in the cytoplasm in activated conventional T-cells whereas predominantly expressed in the nucleus in regulatory T- cells (Treg). The 41 kDa form derived by proteolytic processing is found exclusively in the chromatin fraction of activated Treg cells (By similarity). {ECO:0000250|UniProtKB:Q99JB6, ECO:0000269|PubMed:22678915}

Phospho-FOXP3(S418) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Phospho-FOXP3(S418) Antibody - Images



Dot blot analysis of anti-Phospho-FOXP3-S418 Phospho-specific Pab (Cat. #AP3731a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.5ug per ml.

Phospho-FOXP3(S418) Antibody - Background

The protein is a member of the forkhead/winged-helix family of transcriptional regulators. Defects in this gene are the cause of immunodeficiency polyendocrinopathy, enteropathy, X-linked syndrome (IPEX), also known as X-linked autoimmunity-immunodeficiency syndrome.

Phospho-FOXP3(S418) Antibody - References

Davila, S., et al. Genes Immun. 11(3):232-238(2010) Ganguly, S., et al. J. Invest. Dermatol. 130(4):1013-1022(2010) Jung, D.J., et al. J. Biol. Chem. 285(11):7995-8002(2010) **Phospho-FOXP3(S418) Antibody - Citations**

- <u>AKT1/FOXP3 axis-mediated expression of CerS6 promotes p53 mutant pancreatic</u> tumorigenesis
- Anaphylatoxins Activate Ca, Akt/PI3-Kinase, and FOXO1/FoxP3 in the Retinal Pigment Epithelium.
- <u>Phosphorylation of FOXP3 controls regulatory T cell function and is inhibited by TNF-α in</u> <u>rheumatoid arthritis.</u>