

Phospho-mouse JUN(T289) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3783b

Specification

Phospho-mouse JUN(T289) Antibody - Product Information

Application DB,E
Primary Accession P05627

Other Accession <u>P17325</u>, <u>P56432</u>, <u>P05412</u>, <u>P18870</u>, <u>O77627</u>,

NP 002219.1

Reactivity Mouse

Predicted Bovine, Chicken, Human, Pig, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 35944

Phospho-mouse JUN(T289) Antibody - Additional Information

Gene ID 16476

Other Names

Transcription factor AP-1, AH119, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, Jun A, Jun

Target/Specificity

This mouse JUN Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding T289 of mouse JUN.

Dilution

DB~~1:500

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-mouse JUN(T289) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-mouse JUN(T289) Antibody - Protein Information

Name Jun



Function Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:14707112). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription factor complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (PubMed:2498083). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (By similarity). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS- mediated transcriptional activation of USP28 (By similarity). Binds to the USP28 promoter (By similarity).

Cellular Location


Nucleus {ECO:0000250|UniProtKB:P05412}.

Phospho-mouse JUN(T289) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Phospho-mouse JUN(T289) Antibody - Images



Dot blot analysis of Phospho-mouse JUN-T289 Antibody Phospho-specific Pab (Cat. #AP3783b) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6ug per ml.

Phospho-mouse JUN(T289) Antibody - Background

This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. This gene is





intronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.

Phospho-mouse JUN(T289) Antibody - References

Gonsalves, C., et al. J. Immunol. 185(10):6253-6264(2010) Bozec, A., et al. J. Cell Biol. 190(6):1093-1106(2010) Machida, K., et al. Hepatology 52(2):480-492(2010) Madi, A., et al. BMC Microbiol. 10, 215 (2010) : Johnatty, S.E., et al. PLoS Genet. 6 (7), E1001016 (2010) :