

Phospho-mouse IKKB(S701) Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3851a

Specification

Phospho-mouse IKKB(S701) Antibody - Product Information

Application	DB,E
Primary Accession	O88351
Other Accession	NP_034676.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	86690

Phospho-mouse IKKB(S701) Antibody - Additional Information

Gene ID 16150

Other Names

Inhibitor of nuclear factor kappa-B kinase subunit beta, I-kappa-B-kinase beta, IKK-B, IKK-beta, IkbKB, I-kappa-B kinase 2, IKK2, Nuclear factor NF-kappa-B inhibitor kinase beta, NFKBIKB, Ikbkb, Ikkb

Target/Specificity

This mouse IKKB Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S701 of mouse IKKB.

Dilution

DB~~1:500

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-mouse IKKB(S701) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-mouse IKKB(S701) Antibody - Protein Information

Name Ikbkb

Synonyms Ikkb

Function Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (By similarity). Acts as a part of the canonical IKK complex in the conventional pathway of NF-kappa-B activation (By similarity). Phosphorylates inhibitors of NF-kappa-B on 2 critical serine residues (By similarity). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (By similarity). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (By similarity). In addition to the NF-kappa-B inhibitors, phosphorylates several other components of the signaling pathway including NEMO/IKBKG, NF-kappa-B subunits RELA and NFKB1, as well as IKK-related kinases TBK1 and IKBKE (By similarity). IKK-related kinase phosphorylations may prevent the overproduction of inflammatory mediators since they exert a negative regulation on canonical IKKs (By similarity). Phosphorylates FOXO3, mediating the TNF-dependent inactivation of this pro-apoptotic transcription factor (By similarity). Also phosphorylates other substrates including NAA10, NCOA3, BCL10 and IRS1 (By similarity). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF- mediated RIPK1-dependent cell death (PubMed:[30988283](#)). Phosphorylates the C-terminus of IRF5, stimulating IRF5 homodimerization and translocation into the nucleus (PubMed:[25326420](#)). Following bacterial lipopolysaccharide (LPS)-induced TLR4 endocytosis, phosphorylates STAT1 at 'Thr-748' which restricts interferon signaling and anti-inflammatory responses and promotes innate inflammatory responses (PubMed:[38621137](#)). IKBKB-mediated phosphorylation of STAT1 at 'Thr-748' promotes binding of STAT1 to the ARID5A promoter, resulting in transcriptional activation of ARID5A and subsequent ARID5A-mediated stabilization of IL6 (By similarity). It also promotes binding of STAT1 to the IL12B promoter and activation of IL12B transcription (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:O14920}. Nucleus {ECO:0000250|UniProtKB:O14920}. Membrane raft {ECO:0000250|UniProtKB:O14920}. Note=Colocalized with DPP4 in membrane rafts. {ECO:0000250|UniProtKB:O14920}

Tissue Location


Detected in heart (at protein level) (PubMed:23090968). Expressed in liver, kidney and spleen

Phospho-mouse IKKB(S701) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-mouse IKKB(S701) Antibody - Images



Phospho-mouse IKKB(S701) Antibody - Background

NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA (MIM 164014), or RELB (MIM 604758) to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008, or NFKBIB, MIM 604495), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).

Phospho-mouse IKKB(S701) Antibody - References

Busuttil, V., et al. Proc. Natl. Acad. Sci. U.S.A. 107(42):18061-18066(2010)
Kenneth, N.S., et al. EMBO J. 29(17):2966-2978(2010)
Tsuchiya, Y., et al. Mol. Cell 39(4):570-582(2010)
Farlik, M., et al. Immunity 33(1):25-34(2010)
Dong, X., et al. PLoS Pathog. 6 (7), E1001001 (2010) :