

Phospho-PPP1CA (Thr320) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP3912a**Specification**

Phospho-PPP1CA (Thr320) Antibody - Product Information

Application	WB,E
Primary Accession	P62136
Other Accession	Q3T0E7 , Q8WMS6 , P62137 , P62139 , P62138
Reactivity	Human
Predicted	Bovine, Mouse, Rabbit, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	37512

Phospho-PPP1CA (Thr320) Antibody - Additional Information**Gene ID** 5499**Other Names**

Serine/threonine-protein phosphatase PP1-alpha catalytic subunit, PP-1A, 3.1.3.16, PPP1CA, PPP1A

Target/Specificity

This Phospho-PPP1CA (Thr320) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 291-324 amino acids from the human region of human PPP1CA.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-PPP1CA (Thr320) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-PPP1CA (Thr320) Antibody - Protein Information**Name** PPP1CA

Synonyms PPP1A

Function Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets (PubMed:[28216226](#), PubMed:[30158517](#), PubMed:[35768504](#), PubMed:[35830882](#), PubMed:[35831509](#), PubMed:[36175670](#), PubMed:[39603239](#), PubMed:[39603240](#)). Protein phosphatase 1 (PP1) is essential for cell division, transcription elongation, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis (PubMed:[35768504](#), PubMed:[35830882](#), PubMed:[35831509](#), PubMed:[36175670](#), PubMed:[39603239](#), PubMed:[39603240](#)). Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Catalytic component of the PNUTS-PP1 protein phosphatase complex, a protein phosphatase 1 (PP1) complex that promotes RNA polymerase II transcription pause-release, allowing transcription elongation: the PNUTS-PP1 complex mediates the release of RNA polymerase II from promoter-proximal region of genes by catalyzing dephosphorylation of proteins involved in transcription, such as AFF4, CDK9, MEPCE, INTS12, NCBP1, POLR2M/GDOWN1 and SUPT6H (PubMed:[39603239](#), PubMed:[39603240](#)). The PNUTS-PP1 complex also regulates transcription termination by mediating dephosphorylation of SUPT5H in termination zones downstream of poly(A) sites, thereby promoting deceleration of RNA polymerase II transcription (PubMed:[31677974](#)). PNUTS-PP1 complex is also involved in the response to replication stress by mediating dephosphorylation of POLR2A at 'Ser-5' of the CTD, promoting RNA polymerase II degradation (PubMed:[33264625](#)). PNUTS-PP1 also plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase (PubMed:[20516061](#)). Regulates NEK2 function in terms of kinase activity and centrosome number and splitting, both in the presence and absence of radiation- induced DNA damage (PubMed:[17283141](#)). Regulator of neural tube and optic fissure closure, and enteric neural crest cell (ENCCs) migration during development (By similarity). In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation (PubMed:[21712997](#)). May dephosphorylate CSNK1D and CSNK1E (PubMed:[21712997](#)). Dephosphorylates the 'Ser-418' residue of FOXP3 in regulatory T-cells (Treg) from patients with rheumatoid arthritis, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:[23396208](#)). Dephosphorylates CENPA (PubMed:[25556658](#)). Dephosphorylates the 'Ser-139' residue of ATG16L1 causing dissociation of ATG12-ATG5-ATG16L1 complex, thereby inhibiting autophagy (PubMed:[26083323](#)). Together with PPP1CC (PP1-gamma subunit), dephosphorylates IFIH1/MDA5 and RIG-I leading to their activation and a functional innate immune response (PubMed:[23499489](#)). Core component of the SHOC2-MRAS-PP1c (SMP) holophosphatase complex that regulates the MAPK pathway activation (PubMed:[35768504](#), PubMed:[35830882](#), PubMed:[35831509](#), PubMed:[36175670](#)). The SMP complex specifically dephosphorylates the inhibitory phosphorylation at 'Ser-259' of RAF1 kinase, 'Ser-365' of BRAF kinase and 'Ser-214' of ARAF kinase, stimulating their kinase activities (PubMed:[35768504](#), PubMed:[35830882](#), PubMed:[35831509](#), PubMed:[36175670](#)). The SMP complex enhances the dephosphorylation activity and substrate specificity of PP1c (PubMed:[35768504](#), PubMed:[36175670](#)).

Cellular Location

Cytoplasm. Nucleus. Nucleus, nucleoplasm. Nucleus, nucleolus Note=Primarily nuclear and largely excluded from the nucleolus. Highly mobile in cells and can be relocalized through interaction with targeting subunits. NOM1 plays a role in targeting this protein to the nucleolus. In the presence of PPP1R8 relocalizes from the nucleus to nuclear speckles. Shuttles toward the cytosol during infection with VEEV (PubMed:[29769351](#)).


Phospho-PPP1CA (Thr320) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-PPP1CA (Thr320) Antibody - Images



Western blot analysis of lysates from HeLa cell line, untreated or treated with paclitaxel, 100nM, 20hrs, using Phospho-PPP1CA (Thr320) Antibody (upper) or Beta-actin (lower).

Phospho-PPP1CA (Thr320) Antibody - Background

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. Regulates NEK2 function in terms of kinase activity and centrosome number and splitting, both in the presence and absence of radiation-induced DNA damage. Regulator of neural tube and optic fissure closure, and enteric neural crest cell (ENCCs) migration during development. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E.

Phospho-PPP1CA (Thr320) Antibody - References

- Song Q.,et al.Gene 129:291-295(1993).
Durfee T.,et al.Genes Dev. 7:555-569(1993).
Tung L.,et al.Submitted (APR-1991) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.