

ANKFY1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP4704b

Specification

ANKFY1 Antibody (C-term) - Product Information

Application FC, IHC-P, WB,E

Primary Accession
Reactivity
Host
Clonality
Isotype
Antigen Region

Q9P2R3
Human
Rabbit
Polyclonal
Rabbit IgG
436-465

ANKFY1 Antibody (C-term) - Additional Information

Gene ID 51479

Other Names

Rabankyrin-5, Rank-5, Ankyrin repeat and FYVE domain-containing protein 1, Ankyrin repeats hooked to a zinc finger motif, ANKFY1, ANKHZN, KIAA1255

Target/Specificity

This ANKFY1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 436-465 amino acids from the C-terminal region of human ANKFY1.

Dilution

FC~~1:10~50 IHC-P~~1:50~100 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ANKFY1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ANKFY1 Antibody (C-term) - Protein Information

Name ANKFY1



Synonyms ANKHZN, KIAA1255

Function Proposed effector of Rab5. Binds to phosphatidylinositol 3- phosphate (PI(3)P). Involved in homotypic early endosome fusion and to a lesser extent in heterotypic fusion of chlathrin-coated vesicles with early endosomes. Involved in macropinocytosis; the function is dependent on Rab5-GTP. Required for correct endosomal localization. Involved in the internalization and trafficking of activated tyrosine kinase receptors such as PDGFRB. Regulates the subcellular localization of the retromer complex in a EHD1-dependent manner. Involved in endosome-to-Golgi transport and biosynthetic transport to late endosomes and lysosomes indicative for a regulation of retromer complex-mediated retrograde transport.

Cellular Location

Cytoplasm. Endosome membrane; Peripheral membrane protein. Early endosome. Note=Also associated with endosomal membranes. Localizes to macropinosomes.

Tissue Location


High expression in whole adult brain and intermediate expression in all other tissues and specific brain regions examined, including fetal brain.

ANKFY1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ANKFY1 Antibody (C-term) - Images




ANKFY1 Antibody (C-term) (Cat. #AP4704b) western blot analysis in MDA-MB435,CEM,MCF-7,Jurkat,Hela cell line lysates (35ug/lane).This demonstrates the ANKFY1 antibody detected the ANKFY1 protein (arrow).





ANKFY1 Antibody (C-term) (Cat. #AP4704b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ANKFY1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



ANKFY1 Antibody (C-term) (Cat. #AP4704b) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ANKFY1 Antibody (C-term) - Background

ANKFY1 encodes a cytoplasmic protein that contains a coiled-coil structure and a BTB/POZ domain at its N-terminus, ankyrin repeats in the middle portion, and a FYVE-finger motif at its C-terminus. This protein belongs to a subgroup of double zinc finger proteins which may be involved in vesicle or protein transport.

ANKFY1 Antibody (C-term) - References

Bouslam, N., et al. Hum. Genet. 121 (3-4), 413-420 (2007) Schnatwinkel, C., et al. PLoS Biol. 2 (9), E261 (2004)

ANKFY1 Antibody (C-term) - Citations

- Rapid degradation of progressive ankylosis protein (ANKH) in craniometaphyseal dysplasia.
- Cellular uptake of extracellular vesicles is mediated by clathrin-independent endocytosis and macropinocytosis.