

**CNGA2 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP4858A**

### Specification

#### CNGA2 Antibody (N-term) - Product Information

Application	FC, WB,E
Primary Accession	<a href="#">Q16280</a>
Other Accession	<a href="#">Q00195</a> , <a href="#">Q28718</a> , <a href="#">Q62398</a> , <a href="#">Q03041</a>
Reactivity	Human
Predicted	Bovine, Mouse, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	76048
Antigen Region	75-102

#### CNGA2 Antibody (N-term) - Additional Information

##### Gene ID 1260

##### Other Names

Cyclic nucleotide-gated olfactory channel, Cyclic nucleotide-gated cation channel 2, Cyclic nucleotide-gated channel alpha-2, CNG channel alpha-2, CNG-2, CNG2, CNGA2, CNCA, CNCA1, CNCG2

##### Target/Specificity

This CNGA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 75-102 amino acids from the N-terminal region of human CNGA2.

##### Dilution

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

CNGA2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### CNGA2 Antibody (N-term) - Protein Information

Name CNGA2 {ECO:0000303|PubMed:11764791, ECO:0000312|HGNC:HGNC:2149}

**Function** Pore-forming subunit of the olfactory cyclic nucleotide-gated channel. Operates in the cilia of olfactory sensory neurons where chemical stimulation of the odorant is converted to an electrical signal. Mediates odorant-induced cAMP-dependent Ca(2+) influx triggering neuron depolarization. The rise of intracellular Ca(2+) levels potentiates the olfactory response by activating Ca(2+)- dependent Cl(-) channels, but it also serves as a negative feedback signal to desensitize the channel for rapid adaptation to odorants. Conducts cAMP- and cGMP-gated ion currents, with permeability for monovalent and divalent cations.

#### Cellular Location


Cell projection, cilium membrane {ECO:0000250|UniProtKB:Q00195}; Multi-pass membrane protein

#### CNGA2 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CNGA2 Antibody (N-term) - Images



Western blot analysis of CNGA2 Antibody (N-term) (Cat. #AP4858a) in K562 cell line lysates (35ug/lane). CNGA2 (arrow) was detected using the purified Pab.



CNGA2 Antibody (N-term) (Cat. #AP4858a) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### CNGA2 Antibody (N-term) - Background

CNGA2 represents the alpha subunit of a cyclic nucleotide-gated olfactory channel. The encoded protein contains a carboxy-terminal leucine zipper that mediates channel formation.

#### CNGA2 Antibody (N-term) - References

- Qu, W., et al. J. Gen. Physiol. 127(4):375-389(2006)
- Hofmann, F., et al. Pharmacol. Rev. 57(4):455-462(2005)
- Yoo, D., et al. J. Biol. Chem. 279(8):6863-6873(2004)