

FAS Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50018**Specification**

FAS Antibody - Product Information

Application	WB, IHC
Primary Accession	P25445
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38 KDa
Antigen Region	301-333

FAS Antibody - Additional Information**Gene ID** 355**Other Names**

Tumor necrosis factor receptor superfamily member 6, Apo-1 antigen, Apoptosis-mediating surface antigen FAS, FASLG receptor, CD95, FAS, APT1, FAS1, TNFRSF6

Dilution

WB~~ 1:1000

IHC~~1:50-1:100

Format

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

FAS Antibody - Protein Information**Name** FAS**Synonyms** APT1, FAS1, TNFRSF6**Function**

Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase CASP8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs CASP8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen- stimulated suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Membrane raft [Isoform 3]: Secreted. [Isoform 5]: Secreted.

Tissue Location


Isoform 1 and isoform 6 are expressed at equal levels in resting peripheral blood mononuclear cells. After activation there is an increase in isoform 1 and decrease in the levels of isoform 6.

FAS Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FAS Antibody - Images



Western blot analysis of lysate from HT-1080 cell line, using FAS Antibody (C0355). C0355 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue using Fas antibody.

FAS Antibody - Background

Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro).

FAS Antibody - References

- Itoh N., et al. Cell 66:233-243(1991).
- Oehm A., et al. J. Biol. Chem. 267:10709-10715(1992).
- Liu C., et al. Biochem. J. 310:957-963(1995).
- Cascino I., et al. J. Immunol. 154:2706-2713(1995).
- Cascino I., et al. J. Immunol. 156:13-17(1996).