

CCD92 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5018C**Specification**

CCD92 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q53HC0
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36961
Antigen Region	177-205

CCD92 Antibody (Center) - Additional Information**Gene ID** 80212**Other Names**

Coiled-coil domain-containing protein 92, Limkain beta-2, CCDC92

Target/Specificity

This CCD92 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 177-205 amino acids from the Central region of human CCD92.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCD92 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CCD92 Antibody (Center) - Protein Information**Name** CCDC92

Function Interferon-stimulated protein that plays a role in innate immunity. Strongly inhibits ebolavirus transcription and replication. Forms a complex with viral RNA-bound nucleocapsid NP

and thereby prevents the transport of NP to the cell surface.

Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm

CCD92 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCD92 Antibody (Center) - Images



Western blot analysis of CCD92 Antibody (Center) (Cat. #AP5018c) in mouse testis tissue lysates (35ug/lane). CCD92 (arrow) was detected using the purified Pab.



CCD92 Antibody (Center) (Cat. #AP5018c) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CCD92 Antibody (Center) for

immunohistochemistry. Clinical relevance has not been evaluated.

CCD92 Antibody (Center) - References

Chasman, D.I., et al. PLoS Genet. 5 (11), E1000730 (2009)