

PDLIM1 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50197

Specification

PDLIM1 Antibody - Product Information

Application WB, IF, IHC
Primary Accession O00151

Reactivity
Host
Clonality
Human, Mouse, Rat
Rabbit
Polyclonal

Clonality Polyclor
Calculated MW 36 KDa
Antigen Region 91-119

PDLIM1 Antibody - Additional Information

Gene ID 9124

Other Names

PDZ and LIM domain protein 1, C-terminal LIM domain protein 1, Elfin, LIM domain protein CLP-36, PDLIM1, CLIM1, CLP36

Dilution

WB~~ 1:1000 IF~~1:100 IHC~~1:50-1:100

Format

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

PDLIM1 Antibody - Protein Information

Name PDLIM1

Synonyms CLIM1, CLP36

Function

Cytoskeletal protein that may act as an adapter that brings other proteins (like kinases) to the cytoskeleton (PubMed:10861853). Involved in assembly, disassembly and directioning of stress fibers in fibroblasts. Required for the localization of ACTN1 and PALLD to stress fibers. Required for cell migration and in maintaining cell polarity of fibroblasts (By similarity).

Cellular Location



Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, myofibril, sarcomere, Z line Note=Associates with actin stress fibers

Tissue Location


Strongly expressed in the heart and skeletal muscle, moderately expressed in the spleen, small intestine, colon, placenta, and lung. A lower level expression is seen in liver, thymus, kidney, prostate and pancreas and is not found in the brain, testis, ovary, and peripheral blood leukocytes

PDLIM1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PDLIM1 Antibody - Images




Immunofluorescence analysis of HepG2 cells, using PDLIM1 antibody.



Western blot analysis of lysates from Jurkat, HepG2 cell line (from left to right), using PDLIM1 Antibody(C10012). C10012 was diluted at 1:1000 at each lane. A goat anti-rabbit lgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Tel: 858.875.1900 Fax: 858.875.1999



Immunohistochemistry analysis of paraffin-embedded human heart tissue using PDLIM1 antibody.

PDLIM1 Antibody - Background

Cytoskeletal protein that may act as an adapter that brings other proteins (like kinases) to the cytoskeleton.

PDLIM1 Antibody - References

Kotaka M., et al.J. Cell. Biochem. 72:279-285(1999). Bauer K., et al. Blood 96:4236-4245(2000). Ota T., et al. Nat. Genet. 36:40-45(2004). Deloukas P., et al. Nature 429:375-381(2004). Gevaert K., et al. Nat. Biotechnol. 21:566-569(2003).