

**CHST10 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP50588****Specification**

---

**CHST10 Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">O43529</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42 KDa
Antigen Region	211-239

**CHST10 Antibody - Additional Information****Gene ID** 9486**Other Names**

Carbohydrate sulfotransferase 10, 282-, HNK-1 sulfotransferase, HNK-1ST, HNK1ST, HuHNK-1ST, CHST10

**Dilution**

WB~~ 1:1000

IHC~~1:50-1:100

**Format**Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.**Storage Conditions**

-20°C

**CHST10 Antibody - Protein Information****Name** CHST10 {ECO:0000303|PubMed:23269668, ECO:0000312|HGNC:HGNC:19650}**Function**

Catalyzes the transfer of sulfate from 3'-phosphoadenylyl sulfate (PAPS) to position 3 of terminal glucuronic acid of both protein- and lipid-linked oligosaccharides. Participates in biosynthesis of HNK-1 carbohydrate structure 3-O-sulfo-beta-D-GlcA- (1->3)-beta-D-Gal-(1->4)-D-GlcNAc-R, a sulfated glucuronyl-lactosaminyl residue carried by many neural recognition molecules, which is involved in cell interactions during ontogenetic development and in synaptic plasticity in the adult. May be indirectly involved in synapse plasticity of the hippocampus, via its role in HNK-1 biosynthesis (PubMed: <http://www.uniprot.org/citations/9478973> target="\_blank">9478973</a>). Sulfates terminal glucuronyl residue of the laminin globular (LG)-domain binding epitope on DAG1/alpha-dystroglycan and prevents further polymerization by LARGE1 glycosyltransferase. Likely defines the chain length of LG epitope, conferring binding

specificity to extracellular matrix components (PubMed:<a href="http://www.uniprot.org/citations/32149355" target="\_blank">32149355</a>). Plays a role in down-regulating the steroid hormones. Sulfates glucuronidated estrogens and androgens with an impact in hormone cycle and fertility. Has a preference for glucuronyl moiety at the 3-hydroxyl group of a sterol ring rather than the 17-hydroxyl group, showing high catalytic efficiency for 17beta-estradiol 3-O-(beta-D-glucuronate) and dehydroepiandrosterone 3-O-(beta-D-glucuronate) hormones (PubMed:<a href="http://www.uniprot.org/citations/23269668" target="\_blank">23269668</a>).

#### Cellular Location

Golgi apparatus membrane {ECO:0000250|UniProtKB:O54702}; Single-pass type II membrane protein

#### Tissue Location


In fetal tissues, it is predominantly expressed in brain, and weakly expressed in lung, kidney and liver. In adult, it is highly expressed in brain, testis, ovary, expressed at intermediate level in heart, pancreas, skeletal muscle, spleen and thymus, and weakly expressed in other tissues. In brain, it is expressed at higher level in the frontal lobe.

### CHST10 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CHST10 Antibody - Images



Western blot analysis of lysate from human brain tissue lysate, using CHST10 Antibody (AP50588). AP50588 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using CHST10 antibody.

#### **CHST10 Antibody - Background**

Catalyzes the transfer of sulfate to position 3 of terminal glucuronic acid of both protein- and lipid-linked oligosaccharides. Participates in biosynthesis of HNK-1 carbohydrate structure, a sulfated glucuronyl-lactosaminyl residue carried by many neural recognition molecules, which is involved in cell interactions during ontogenetic development and in synaptic plasticity in the adult. May be indirectly involved in synapse plasticity of the hippocampus, via its role in HNK-1 biosynthesis.

#### **CHST10 Antibody - References**

- Ong E.,et al.J. Biol. Chem. 273:5190-5195(1998).
- Yu W.,et al.Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases.
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Hillier L.W.,et al.Nature 434:724-731(2005).
- Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.