

FGF13 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP50639

Specification

FGF13 Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Antigen Region

WB
O92913
Human, Mouse, Rat
Rabbit
Polyclonal
28,22,29,26 KDa
175-198

FGF13 Antibody - Additional Information

Gene ID 2258

Other Names

Fibroblast growth factor 13, FGF-13, Fibroblast growth factor homologous factor 2, FHF-2, FGF13, FHF2

Dilution

WB~~1:1000

Format

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

FGF13 Antibody - Protein Information


Name FGF13 (HGNC:3670)

Synonyms FHF2

Function

Microtubule-binding protein which directly binds tubulin and is involved in both polymerization and stabilization of microtubules (By similarity). Through its action on microtubules, may participate in the refinement of axons by negatively regulating axonal and leading processes branching (By similarity). Plays a crucial role in neuron polarization and migration in the cerebral cortex and the hippocampus (By similarity). Regulates voltage-gated sodium channel transport and function (PubMed:15282281, PubMed:33245860, PubMed:36696443, Nay also play a role in MAPK signaling (By similarity). Required for the development of axonal







initial segment-targeting inhibitory GABAergic synapses made by chandelier neurons (By similarity).

Cellular Location

[Isoform 1]: Nucleus [Isoform 3]: Cytoplasm. Nucleus [Isoform 5]: Cytoplasm {ECO:0000250|UniProtKB:P61329}. Nucleus {ECO:0000250|UniProtKB:P61329}

Tissue Location


Ubiquitously expressed. Predominantly expressed in the nervous system.

FGF13 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FGF13 Antibody - Images



Western blot analysis of lysate from human heart tissue lysate, using FGF13 Antibody(AP50639). AP50639 was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

FGF13 Antibody - Background

Microtubule-binding protein which directly binds tubulin and is involved in both polymerization and stabilization of microtubules. Through its action on microtubules, may participate to the refinement of axons by negatively regulating axonal and leading processes branching. Plays a crucial role in neuron polarization and migration in the cerebral cortex and the hippocampus. May also play a role in MAPK signaling.





FGF13 Antibody - References

Smallwood P.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:9850-9857(1996). Gecz J.,et al.Hum. Genet. 104:56-63(1999). Ota T.,et al.Nat. Genet. 36:40-45(2004). Ross M.T.,et al.Nature 434:325-337(2005). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.