

**THOC4 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP50789****Specification**

---

**THOC4 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q86V81</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27 KDa
Antigen Region	143-170

**THOC4 Antibody - Additional Information****Gene ID** 10189**Other Names**

THO complex subunit 4, Tho4, Ally of AML-1 and LEF-1, Aly/REF export factor, Transcriptional coactivator Aly/REF, bZIP-enhancing factor BEF, ALYREF, ALY, BEF, THOC4

**Dilution**

WB~~ 1:1000

**Format**

Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

**Storage Conditions**

-20°C

**THOC4 Antibody - Protein Information****Name** ALYREF**Synonyms** ALY, BEF, THOC4**Function**

Export adapter involved in nuclear export of spliced and unspliced mRNA. Binds mRNA which is thought to be transferred to the NXF1-NXT1 heterodimer for export (TAP/NFX1 pathway)  
(PubMed: <a href="http://www.uniprot.org/citations/15833825" target="\_blank">15833825</a>,  
PubMed: <a href="http://www.uniprot.org/citations/15998806" target="\_blank">15998806</a>,  
PubMed: <a href="http://www.uniprot.org/citations/17190602" target="\_blank">17190602</a>,  
PubMed: <a href="http://www.uniprot.org/citations/11707413" target="\_blank">11707413</a>,  
PubMed: <a href="http://www.uniprot.org/citations/11675789" target="\_blank">11675789</a>,  
PubMed: <a href="http://www.uniprot.org/citations/11979277" target="\_blank">11979277</a>,  
PubMed: <a href="http://www.uniprot.org/citations/18364396" target="\_blank">18364396</a>,

PubMed:<a href="http://www.uniprot.org/citations/22144908" target="\_blank">22144908</a>, PubMed:<a href="http://www.uniprot.org/citations/22893130" target="\_blank">22893130</a>, PubMed:<a href="http://www.uniprot.org/citations/23222130" target="\_blank">23222130</a>, PubMed:<a href="http://www.uniprot.org/citations/25662211" target="\_blank">25662211</a>). Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/15833825" target="\_blank">15833825</a>, PubMed:<a href="http://www.uniprot.org/citations/15998806" target="\_blank">15998806</a>, PubMed:<a href="http://www.uniprot.org/citations/17190602" target="\_blank">17190602</a>). TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm (PubMed:<a href="http://www.uniprot.org/citations/15833825" target="\_blank">15833825</a>, PubMed:<a href="http://www.uniprot.org/citations/15998806" target="\_blank">15998806</a>, PubMed:<a href="http://www.uniprot.org/citations/17190602" target="\_blank">17190602</a>). TREX recruitment occurs via an interaction between ALYREF/THOC4 and the cap-binding protein NCBP1 (PubMed:<a href="http://www.uniprot.org/citations/15833825" target="\_blank">15833825</a>, PubMed:<a href="http://www.uniprot.org/citations/15998806" target="\_blank">15998806</a>, PubMed:<a href="http://www.uniprot.org/citations/17190602" target="\_blank">17190602</a>). The TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV) intronless mRNAs and infectious virus production; ALYREF/THOC4 mediates the recruitment of the TREX complex to the intronless viral mRNA (PubMed:<a href="http://www.uniprot.org/citations/18974867" target="\_blank">18974867</a>). Required for TREX complex assembly and for linking DDX39B to the cap-binding complex (CBC) (PubMed:<a href="http://www.uniprot.org/citations/15998806" target="\_blank">15998806</a>, PubMed:<a href="http://www.uniprot.org/citations/17984224" target="\_blank">17984224</a>). In conjunction with THOC5 functions in NXF1-NXT1 mediated nuclear export of HSP70 mRNA; both proteins enhance the RNA binding activity of NXF1 and are required for NXF1 localization to the nuclear rim (PubMed:<a href="http://www.uniprot.org/citations/19165146" target="\_blank">19165146</a>). Involved in the nuclear export of intronless mRNA; proposed to be recruited to intronless mRNA by ATP-bound DDX39B. Involved in transcription elongation and genome stability (PubMed:<a href="http://www.uniprot.org/citations/12438613" target="\_blank">12438613</a>, PubMed:<a href="http://www.uniprot.org/citations/17984224" target="\_blank">17984224</a>). Involved in mRNA export of C5-methylcytosine (m5C)-containing mRNAs: specifically recognizes and binds m5C mRNAs and mediates their nucleo- cytoplasmic shuttling (PubMed:<a href="http://www.uniprot.org/citations/28418038" target="\_blank">28418038</a>).

### Cellular Location

Nucleus. Nucleus speckle Cytoplasm Note=Colocalizes with the core EJC, ALYREF/THOC4, NXF1 and DDX39B in the nucleus and nuclear speckles. Travels to the cytoplasm as part of the exon junction complex (EJC) bound to mRNA (PubMed:19324961) Localizes to regions surrounding nuclear speckles known as perispeckles in which TREX complex assembly seems to occur (PubMed:23826332)

### Tissue Location

Expressed in a wide variety of cancer types.


### THOC4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### THOC4 Antibody - Images



Western blot analysis of lysate from KG-1 cell line, using THOC4 Antibody (AP50789). AP50789 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35 µg.

### THOC4 Antibody - Background

Export adapter involved in nuclear export of spliced and unspliced mRNA. Binds mRNA which is thought to be transferred to the NXF1-NXT1 heterodimer for export (TAP/NFX1 pathway). Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm. TREX recruitment occurs via an interaction between ALYREF/THOC4 and the cap-binding protein NCBP1. The TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV) intronless mRNAs and infectious virus production; ALYREF/THOC4 mediates the recruitment of the TREX complex to the intronless viral mRNA. Required for TREX complex assembly and for linking DDX39B to the cap-binding complex (CBC). In conjunction with THOC5 functions in NXF1-NXT1 mediated nuclear export of HSP70 mRNA; both proteins enhance the RNA binding activity of NXF1 and are required for NXF1 localization to the nuclear rim. Involved in the nuclear export of intronless mRNA; proposed to be recruited to intronless mRNA by ATP-bound DDX39B. Involved in transcription elongation and genome stability.

### THOC4 Antibody - References

Zody M.C., et al. Nature 440:1045-1049 (2006).  
 Quadroni M., et al. Submitted (OCT-2004) to UniProtKB.  
 Bienvenut W.V., et al. Submitted (DEC-2008) to UniProtKB.  
 Wichmann I., et al. Hum. Immunol. 60:57-62 (1999).

Rappsilber J., et al. Genome Res. 12:1231-1245(2002).