

#### **Cerebellin 4 Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51048

#### **Specification**

#### **Cerebellin 4 Antibody - Product Information**

Application WB
Primary Accession O9NTU7

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 22 KDa
Antigen Region 131 - 190

### **Cerebellin 4 Antibody - Additional Information**

**Gene ID 140689** 

**Other Names** 

Cerebellin-4, Cerebellin-like glycoprotein 1, CBLN4, CBLNL1

Target/Specificity

KLH conjugated synthetic peptide derived from human Cerebellin 4

**Dilution** 

WB~~ 1:1000

**Format** 

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage** 

Store at -20 °C.Stable for 12 months from date of receipt

# **Cerebellin 4 Antibody - Protein Information**

Name CBLN4

Synonyms CBLNL1

#### **Function**

Acts as a synaptic organizer in specific subsets of neurons in the brain (By similarity). Essential for the formation and maintenance of inhibitory GABAergic synapses (By similarity). Promotes the development of dendrite-targeting inhibitory GABAergic synapses made by somatostatin-positive interneurons (By similarity). May contribute to the function of ventral medial habenula region of the brain implicated in the regulation of anxiety-related behaviors (By similarity). May play a role in CBLN3 export from the endoplasmic reticulum and secretion (By similarity).

### **Cellular Location**




Secreted {ECO:0000250|UniProtKB:Q8BME9}. Synapse {ECO:0000250|UniProtKB:Q8BME9}. Note=Detected at GABAergic synapses {ECO:0000250|UniProtKB:Q8BME9}

# **Cerebellin 4 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

#### Cerebellin 4 Antibody - Images



All lanes : Anti-Cerebellin 4 Antibody at 1:1000 dilution Lane 1: rat liver lysates Lane 2: 293T/17 whole cell lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### Cerebellin 4 Antibody - Background

May be involved in synaptic functions in the CNS. May play a role in CBLN3 export from the endoplasmic reticulum and secretion (By similarity).

## **Cerebellin 4 Antibody - References**

Clark H.F., et al. Genome Res. 13:2265-2270(2003). Ota T., et al. Nat. Genet. 36:40-45(2004). Deloukas P., et al. Nature 414:865-871(2001). Zhang Z., et al. Protein Sci. 13:2819-2824(2004).