

**Cadherin 23 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51067****Specification**

---

**Cadherin 23 Antibody - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">O9H251</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59 KDa

**Cadherin 23 Antibody - Additional Information****Gene ID** 64072**Other Names**

Cadherin-23, Otocadherin, CDH23, KIAA1774, KIAA1812

**Target/Specificity**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Cadherin 23. The exact sequence is proprietary.

**Dilution**

WB~~1:1000

IHC-P~~N/A

E~~N/A

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Cadherin 23 Antibody - Protein Information****Name** CDH23 {ECO:0000303|PubMed:11138009, ECO:0000312|HGNC:HGNC:13733}**Function**

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells. CDH23 is required for establishing and/or maintaining the proper organization of the stereocilia bundle of hair cells in the cochlea and the vestibule during late embryonic/early postnatal development. It is part of the functional network formed by USH1C, USH1G, CDH23 and MYO7A that mediates mechanotransduction in cochlear hair cells. Required for normal hearing.

**Cellular Location**

Cell membrane; Single-pass type I membrane protein

**Tissue Location**

Particularly strong expression in the retina (PubMed:11138009). Found also in the cochlea

**Cadherin 23 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Cadherin 23 Antibody - Images****Cadherin 23 Antibody - Background**

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells. CDH23 is required for establishing and/or maintaining the proper organization of the stereocilia bundle of hair cells in the cochlea and the vestibule during late embryonic/early postnatal development. It is part of the functional network formed by USH1C, USH1G, CDH23 and MYO7A that mediates mechanotransduction in cochlear hair cells. Required for normal hearing.

**Cadherin 23 Antibody - References**

Bolz H.,et al.Nat. Genet. 27:108-112(2001).  
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).  
Lagziel A.,et al.Dev. Biol. 280:295-306(2005).  
Deloukas P.,et al.Nature 429:375-381(2004).  
Nagase T.,et al.DNA Res. 8:85-95(2001).