

Cytochrome P450 1B1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51142**Specification**

Cytochrome P450 1B1 Antibody - Product Information

Application	WB, E
Primary Accession	Q16678
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70 KDa

Cytochrome P450 1B1 Antibody - Additional Information**Gene ID** 1545**Other Names**

Cytochrome P450 1B1, CYP1B1, CYP1B1

Dilution

WB~~1:1000

E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Cytochrome P450 1B1 Antibody - Protein Information**Name** CYP1B1 {ECO:0000303|PubMed:8910454, ECO:0000312|HGNC:HGNC:2597}**Function**

A cytochrome P450 monooxygenase involved in the metabolism of various endogenous substrates, including fatty acids, steroid hormones and vitamins (PubMed:10681376, PubMed:11555828, PubMed:12865317, PubMed:15258110, PubMed:20972997).

Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:10681376, PubMed:11555828, PubMed:12865317, PubMed:15258110, PubMed:20972997).

[15258110](http://www.uniprot.org/citations/15258110), PubMed: [20972997](http://www.uniprot.org/citations/20972997)). Exhibits catalytic activity for the formation of hydroxyestrogens from estrone (E1) and 17beta-estradiol (E2), namely 2- and 4-hydroxy E1 and E2. Displays a predominant hydroxylase activity toward E2 at the C-4 position (PubMed: [11555828](http://www.uniprot.org/citations/11555828), PubMed: [12865317](http://www.uniprot.org/citations/12865317)). Metabolizes testosterone and progesterone to B or D ring hydroxylated metabolites (PubMed: [10426814](http://www.uniprot.org/citations/10426814)). May act as a major enzyme for all-trans retinoic acid biosynthesis in extrahepatic tissues. Catalyzes two successive oxidative transformation of all-trans retinol to all-trans retinal and then to the active form all-trans retinoic acid (PubMed: [10681376](http://www.uniprot.org/citations/10681376), PubMed: [15258110](http://www.uniprot.org/citations/15258110)). Catalyzes the epoxidation of double bonds of certain PUFA. Converts arachidonic acid toward epoxyeicosatrienoic acid (EpETRe) regioisomers, 8,9-, 11,12-, and 14,15- EpETRe, that function as lipid mediators in the vascular system (PubMed: [20972997](http://www.uniprot.org/citations/20972997)). Additionally, displays dehydratase activity toward oxygenated eicosanoids hydroperoxyeicosatetraenoates (HpETEs). This activity is independent of cytochrome P450 reductase, NADPH, and O₂ (PubMed: [21068195](http://www.uniprot.org/citations/21068195)). Also involved in the oxidative metabolism of xenobiotics, particularly converting polycyclic aromatic hydrocarbons and heterocyclic aryl amines procarcinogens to DNA-damaging products (PubMed: [10426814](http://www.uniprot.org/citations/10426814)). Plays an important role in retinal vascular development. Under hyperoxic O₂ conditions, promotes retinal angiogenesis and capillary morphogenesis, likely by metabolizing the oxygenated products generated during the oxidative stress. Also, contributes to oxidative homeostasis and ultrastructural organization and function of trabecular meshwork tissue through modulation of POSTN expression (By similarity).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q64429}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q64429}. Microsome membrane {ECO:0000250|UniProtKB:Q64429}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q64429}. Mitochondrion {ECO:0000250|UniProtKB:Q64429}. Note=Located primarily in endoplasmic reticulum. Upon treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), CYP1B1 is also targeted to mitochondria {ECO:0000250|UniProtKB:Q64429}

Tissue Location

Expressed in heart, brain, lung, skeletal muscle, kidney, spleen, thymus, prostate, testis, ovary, small intestine, colon, and peripheral blood leukocytes (PubMed:8175734). Expressed in retinal endothelial cells and umbilical vein endothelial cells (at protein level) (PubMed:19005183).

Cytochrome P450 1B1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cytochrome P450 1B1 Antibody - Images**Cytochrome P450 1B1 Antibody - Background**

Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics.

Cytochrome P450 1B1 Antibody - References

Sutter T.R.,et al.J. Biol. Chem. 269:13092-13099(1994).
Tang Y.M.,et al.J. Biol. Chem. 271:28324-28330(1996).
Gorry M.C.,et al.Submitted (NOV-2001) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Guillemette C.,et al.Submitted (JUL-1999) to the EMBL/GenBank/DDBJ databases.