

# Cytochrome P450 2A7 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51145

### Specification

# Cytochrome P450 2A7 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC-P, E <u>P20853</u> Human, Mouse, Rat Rabbit Polyclonal 56 KDa

### Cytochrome P450 2A7 Antibody - Additional Information

Gene ID 1549

**Other Names** Cytochrome P450 2A7, CYPIIA7, Cytochrome P450 IIA4, CYP2A7

Dilution WB~~1:1000 IHC-P~~N/A E~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

#### Cytochrome P450 2A7 Antibody - Protein Information

Name CYP2A7

Function

Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics.

**Cellular Location** Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

#### Cytochrome P450 2A7 Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Cytochrome P450 2A7 Antibody - Images

#### Cytochrome P450 2A7 Antibody - Background

Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics.

#### Cytochrome P450 2A7 Antibody - References

Yamano S., et al. Biochemistry 29:1322-1329(1990). Fernandez-Salguero P., et al. Am. J. Hum. Genet. 57:651-660(1995). Grimwood J., et al. Nature 428:529-535(2004).