

Cytochrome P450 4A11/22 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51150

Specification

Cytochrome P450 4A11/22 Antibody - Product Information

Application	WB, IP, E
Primary Accession	O02928
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	59 KDa

Cytochrome P450 4A11/22 Antibody - Additional Information

Gene ID 1579

Other Names

Cytochrome P450 4A11, 20-hydroxyeicosatetraenoic acid synthase, 20-HETE synthase, CYP4A11, CYP4A11, Cytochrome P-450HK-omega, Cytochrome P450HL-omega, Fatty acid omega-hydroxylase, Lauric acid omega-hydroxylase, CYP4A11, CYP4A2

Dilution

WB~~1:1000
IP~~N/A
E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Cytochrome P450 4A11/22 Antibody - Protein Information

Name CYP4A11 {ECO:0000303|PubMed:8274222, ECO:0000312|HGNC:HGNC:2642}

Function

A cytochrome P450 monooxygenase involved in the metabolism of fatty acids and their oxygenated derivatives (oxylipins) (PubMed: [10553002](http://www.uniprot.org/citations/10553002) target="_blank">10553002, PubMed: [10660572](http://www.uniprot.org/citations/10660572) target="_blank">10660572, PubMed: [15611369](http://www.uniprot.org/citations/15611369) target="_blank">15611369, PubMed: [1739747](http://www.uniprot.org/citations/1739747) target="_blank">1739747, PubMed: [7679927](http://www.uniprot.org/citations/7679927) target="_blank">7679927, PubMed: [8914854](http://www.uniprot.org/citations/8914854) target="_blank">8914854). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed: <a

<http://www.uniprot.org/citations/10553002> target="_blank">10553002, PubMed:10660572, PubMed:15611369, PubMed:1739747, PubMed:7679927, PubMed:8914854). Catalyzes predominantly the oxidation of the terminal carbon (omega-oxidation) of saturated and unsaturated fatty acids, the catalytic efficiency decreasing in the following order: dodecanoic > tetradecanoic > (9Z)-octadecenoic > (9Z,12Z)- octadecadienoic > hexadecanoic acid (PubMed:10553002, PubMed:10660572). Acts as a major omega-hydroxylase for dodecanoic (lauric) acid in liver (PubMed:15611369, PubMed:1739747, PubMed:7679927, PubMed:8914854). Participates in omega-hydroxylation of (5Z,8Z,11Z,14Z)-eicosatetraenoic acid (arachidonate) to 20-hydroxyeicosatetraenoic acid (20-HETE), a signaling molecule acting both as vasoconstrictive and natriuretic with overall effect on arterial blood pressure (PubMed:10620324, PubMed:10660572, PubMed:15611369). Can also catalyze the oxidation of the penultimate carbon (omega-1 oxidation) of fatty acids with lower efficiency (PubMed:7679927). May contribute to the degradation of saturated very long-chain fatty acids (VLCFAs) such as docosanoic acid, by catalyzing successive omega-oxidations to the corresponding dicarboxylic acid, thereby initiating chain shortening (PubMed:18182499). Omega-hydroxylates (9R,10S)-epoxy-octadecanoate stereoisomer (PubMed:15145985). Plays a minor role in omega-oxidation of long-chain 3-hydroxy fatty acids (PubMed:18065749). Has little activity toward prostaglandins A1 and E1 (PubMed:7679927).

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Tissue Location

Expressed in liver (PubMed:7679927). Expressed in S2 and S3 segments of proximal tubules in cortex and outer medulla of kidney (PubMed:10660572, PubMed:7679927).

Cytochrome P450 4A11/22 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cytochrome P450 4A11/22 Antibody - Images**Cytochrome P450 4A11/22 Antibody - Background**

Catalyzes the omega- and (omega-1)-hydroxylation of various fatty acids such as laurate, myristate and palmitate. Has little activity toward prostaglandins A1 and E1. Oxidizes arachidonic acid to 20-hydroxyeicosatetraenoic acid (20-HETE).

Cytochrome P450 4A11/22 Antibody - References

Palmer C.N.A.,et al.Biochim. Biophys. Acta 1172:161-166(1993).
Kawashima H.,et al.J. Biochem. 116:74-80(1994).
Imaoka S.,et al.DNA Cell Biol. 12:893-899(1993).
Bellamine A.,et al.Arch. Biochem. Biophys. 409:221-227(2003).
Gregory S.G.,et al.Nature 441:315-321(2006).