

Ephrin B3 Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51181

Specification

Ephrin B3 Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, ICC, E
O15768
Human, Mouse, Rat
Rabbit
Polyclonal
36 KDa

Ephrin B3 Antibody - Additional Information

Gene ID 1949

Other Names

Ephrin-B3, EPH-related receptor transmembrane ligand ELK-L3, EPH-related receptor tyrosine kinase ligand 8, LERK-8, EFNB3, EPLG8, LERK8

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Ephrin B3 Antibody - Protein Information

Name EFNB3

Synonyms EPLG8, LERK8

Function

Cell surface transmembrane ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. May play a pivotal role in forebrain function. Binds to, and induce the collapse of, commissural axons/growth cones in vitro. May play a role in constraining the orientation of longitudinally projecting axons (By similarity).

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location

Highly expressed in brain; expressed in embryonic floor plate, roof plate and hindbrain segments



Ephrin B3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Ephrin B3 Antibody - Images

Ephrin B3 Antibody - Background

Cell surface transmembrane ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. May play a pivotal role in forebrain function. Binds to, and induce the collapse of, commissural axons/growth cones in vitro. May play a role in constraining the orientation of longitudinally projecting axons (By similarity).

Ephrin B3 Antibody - References

Cerretti D.P.,et al.Submitted (JUL-1996) to the EMBL/GenBank/DDBJ databases. Tang X.X.,et al.Genomics 41:17-24(1997). Gale N.W.,et al.Oncogene 13:1343-1352(1996). Ota T.,et al.Nat. Genet. 36:40-45(2004). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.