

EXO1 Antibody**Purified Rabbit Polyclonal Antibody (Pab)****Catalog # AP51200****Specification****EXO1 Antibody - Product Information**

Application	WB, ICC, E
Primary Accession	Q9UQ84
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	94 KDa

EXO1 Antibody - Additional Information**Gene ID 9156****Other Names**

Exonuclease 1, hExo1, 31--, Exonuclease I, hExo1, EXO1, EXOI, HEX1

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human EXO1. The exact sequence is proprietary.

Dilution

WB~~1:1000

ICC~~N/A

E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

EXO1 Antibody - Protein Information**Name EXO1****Synonyms EXOI, HEX1****Function**

5'→3' double-stranded DNA exonuclease which may also possess a cryptic 3'→5' double-stranded DNA exonuclease activity. Functions in DNA mismatch repair (MMR) to excise mismatch-containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. Also exhibits endonuclease activity against 5'-overhanging flap structures similar to those generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. Required for somatic hypermutation (SHM) and class switch recombination (CSR) of

immunoglobulin genes. Essential for male and female meiosis.

Cellular Location

Nucleus Note=Colocalizes with PCNA to discrete nuclear foci in S-phase

Tissue Location

Highly expressed in bone marrow, testis and thymus. Expressed at lower levels in colon, lymph nodes, ovary, placenta, prostate, small intestine, spleen and stomach

EXO1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EXO1 Antibody - Images

EXO1 Antibody - Background

5'->3' double-stranded DNA exonuclease which may also possess a cryptic 3'->5' double-stranded DNA exonuclease activity. Functions in DNA mismatch repair (MMR) to excise mismatch- containing DNA tracts directed by strand breaks located either 5' or 3' to the mismatch. Also exhibits endonuclease activity against 5'-overhanging flap structures similar to those generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. Required for somatic hypermutation (SHM) and class switch recombination (CSR) of immunoglobulin genes. Essential for male and female meiosis.

EXO1 Antibody - References

Schmutte C.,et al.Cancer Res. 58:4537-4542(1998).
Tishkoff D.X.,et al.Cancer Res. 58:5027-5031(1998).
Wilson D.M. III,et al.Nucleic Acids Res. 26:3762-3768(1998).
Qiu J.,et al.J. Biol. Chem. 274:17893-17900(1999).
Gregory S.G.,et al.Nature 441:315-321(2006).