

Frizzled 6 Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51220

Specification

Frizzled 6 Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IP, E
060353
Human, Mouse, Rat
Rabbit
Polyclonal
80 KDa

Frizzled 6 Antibody - Additional Information

Gene ID 8323

Other Names

Frizzled-6, Fz-6, hFz6, FZD6

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Frizzled 6. The exact sequence is proprietary.

Dilution

WB~~1:1000 IP~~N/A E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Frizzled 6 Antibody - Protein Information

Name FZD6

Function

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues. Together with



FZD3, is involved in the neural tube closure and plays a role in the regulation of the establishment of planar cell polarity (PCP), particularly in the orientation of asymmetric bundles of stereocilia on the apical faces of a subset of auditory and vestibular sensory cells located in the inner ear (By similarity).

Cellular Location

Membrane {ECO:0000250|UniProtKB:Q61089}; Multi- pass membrane protein. Cell membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Cell surface {ECO:0000250|UniProtKB:Q61089}. Apical cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q61089}; Multi-pass membrane protein. Note=Colocalizes with FZD3 at the apical face of cells (By similarity). Localizes to the endoplasmic reticulum membrane in the presence of LMBR1L (By similarity). {ECO:0000250|UniProtKB:Q61089}

Tissue Location

Detected in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, thymus, prostate, testis, ovary, small intestine and colon. In the fetus, expressed in brain, lung, liver and kidney

Frizzled 6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Frizzled 6 Antibody - Images

Frizzled 6 Antibody - Background

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK- 3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.

Frizzled 6 Antibody - References

Tokuhara M.,et al.Biochem. Biophys. Res. Commun. 243:622-627(1998). Gazit A.,et al.Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases. Tanner S.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 98:13901-13906(2001). Suwa M.,et al.Submitted (JUL-2001) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).